

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 23(1) January-April 2025]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology, Pantnagar



ADVISORY BOARD

Patron

Prof. Manmohan Singh Chauhan, Ph.D., Vice-Cancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. A.S. Nain, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. A.H. Ahmad, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.P.S. Gangwar, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Subhash Chandra, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Anil Kumar Gaur, Ph.D., Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A

Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Indu S Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

M.P. Pandey, Ph.D., Ex. Vice Chancellor, BAU, Ranchi & IGKV, Raipur, Director General, IAT, Allahabad, India

Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agri. & Tech., Pantnagar, India

Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

Editor-in-Chief

Manoranjan Dutta, Ph.D., Ex Head, Germplasm Evaluation Division, National Bureau of Plant Genetic Resources, New Delhi, India

Managing Editor

S.N. Tiwari, Ph.D., Professor (Retd.) & Ex-Director Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head

Brijesh Dumka, Developer & Programmer

CONTENTS

<i>In-silico</i> analysis of curcumin conjugates targeting the Wnt signaling pathway in Breast Cancer Stem Cells	1
KANCHAN GAIROLA and SHIV KUMAR DUBEY	
Investigating <i>in vitro</i> direct antagonistic effect of endophytic bacteria against <i>Alternaria brassicicola</i>	8
SHIVANGI KRISHNATRA and A. K SHARMA	
Impact of altitude on photosynthetic and biochemical profile of <i>Didymocarpus pedicellatus</i> R.Br.: an antiurolithiatic Himalayan herb	18
DIVYA and PREETI CHATURVEDI	
Impact of Integrated Nutrient Management (INM) on growth, yield, quality and soil fertility status in sugarcane-ratoon system	25
JYOTI PAWAR and DHEER SINGH	
Mapping and evaluation of soil macronutrient and micronutrient status in Muzaffarnagar district of India	32
RAUSHAN KUMAR and G. R. SINGH	
Study of shift in cropping pattern in northern dry zone of Karnataka	45
ASHWINI HEBBAR and SUMA A. P.	
Changing weather conditions during summer and early monsoon season in the <i>Tarai</i> region of Uttarakhand	51
SHIVANI KOTHIYAL and R.K. SINGH	
Nutrients enhancing flowering characteristics in Mango (<i>Mangifera indica</i> cv. Dashehari) under medium density planting	57
KULDEEP, ASHOK KUMAR SINGH and SHAILESH CHANDRA SHANKHDHAR	
Nutrients and antioxidants potential of star fruit (<i>Averrhoa carambola</i> L.)	66
ABHIMA K. MOORTHY and LAKSHMY P. S.	
Physico-chemical and anti-nutritional properties of predigested composite flour mix from corn and green gram	76
MANISHA RANI and ANJU KUMARI	
Standardisation and quality evaluation of coconut milk yoghurt	84
RINIYA THAJ and LAKSHMY P. S.	
Study on growth performance and morphometric traits of Chaugarkha goat kids in Almora hills of Uttarakhand	93
UMA NAULIA and B. N. SHAHI	

Bacterial isolates from tracheo-bronchial aspirates of healthy and pneumonic cattle ASMITA NARANG, CHARANJIT SINGH, MUDIT CHANDRA and DHIRAJ KUMAR GUPTA	99
Successful management of notoedric mange in two domestic cats: A case report ASMITA NARANG, GURPREET SINGH PREET, JASNIT SINGH and HARKIRAT SINGH	106
Dietary supplementation of formulated fish-specific mineral mixtures improved the growth, nutrient composition and health status of <i>Cyprinus carpio</i> fingerlings ABHED PANDEY, UDEYBIR SINGH CHAHAL and ANJU VIJAYAN	110
Impact of deep cryogenic treatment on microstructural and electrical properties of recycled aluminium alloys BIRENDRA SINGH KARKI and ANADI MISRA	118
Assessing farmers' attitudes and factors influencing livelihood diversification in Nainital District of Uttarakhand NEHA PANDEY, AMARDEEP and V.L.V. KAMESWARI	126
Perceived Benefits of Tribal Sub Plan (TSP) Project on tribal beneficiaries in Udham Singh Nagar District of Uttarakhand ARPITA SHARMA KANDPAL, JITENDRA KWATRA, VLV KAMESWARI and AMARDEEP	133

Impact of altitude on photosynthetic and biochemical profile of *Didymocarpus pedicellatus* R.Br.: an antiurolithiatic Himalayan herb

DIVYA and PREETI CHATURVEDI*

Department of Biological Sciences, College of Basic Sciences and Humanities, G. B. Pant University of Agriculture and Technology, Pantnagar-263145 (U.S. Nagar, Uttarakhand)

*Corresponding author's email id: pchaturvedi.biolsc@gmail.com

ABSTRACT: Altitude plays an important role in shaping plant's physiological and biochemical behavior. This ultimately helps the plant in acclimatization and adaptation in varying habitats spread across different altitudinal gradients. The current work focuses to study the biochemical and physiological profile of *D. pedicellatus* across varying elevations (700, 1186, 1370 and 1640 m). Total polyphenolic content including phenols, flavonoids and tannins were quantitatively estimated using standard protocols. Proteins, photosynthetic pigments and macronutrients were evaluated using standard available protocols with mild modifications. Data was statistically analyzed and means were compared for significance at $p < 0.05$. Results revealed statistically significant variation in all the studied parameters except for phosphorous content. Total phenol (23.26 %), tannin (63.75 %) and protein (175.7 %) content increased effectively whereas total flavonoids decreased (84.84 %) with rise in elevation. Photosynthetic pigments like chlorophyll a, b and carotenoids also followed the decreasing trend by 36.46 %, 39.98 % and 36.24 % respectively, with increasing altitude. Macronutrients like nitrogen, potassium and phosphorus also declined by 55.43 %, 57.67 % and 20.71 % respectively. The decline in photosynthetic pigments and macronutrient content highlights the decrease in photosynthetic and mineral uptake capacity of *D. pedicellatus* at higher altitudes. The study suggests that A4 populations (growing at 1640 m) are richer in polyphenolic compounds and hence more suitable for medicinal purposes. Hence, for bringing the plant in cultivation, A4 populations may be selected as elite planting material in order to get optimum yield of medicinally important compounds.

Keywords: Altitudinal variation, environmental stress, medicinal, nutrients, polyphenols, sustainability

Large number of environmental factors influence plant growth, development and its physiological and biochemical responses. Physical factors like altitude affect solar radiations, temperature, precipitation, atmospheric pressure and nutrient availability (Körner, 2007; Du *et al.*, 2017). Elevation shift leads to notable changes in plant morphology and metabolism, driving the development of altitude-specific mechanism to ensure plant survival (Zhao *et al.*, 2020; Dongdong *et al.*, 2020). *Didymocarpus pedicellatus* (Gesneriaceae) is an endemic Himalayan herb used for curing kidney stones accompanied by nephroprotective, antibacterial, antioxidant and anti-inflammatory properties (Goyal *et al.*, 2015). The plant is distributed in the Indian Himalayan region at an altitude ranging from 700 m to 2000 m inhabiting moist shady places on rocks

mostly near a river stream. Leaves and roots of this plant contain various phytochemical compounds like phenols, quinochalones, chalcones, flavanines, terpenes, fatty acids and steroids (Nanjala *et al.*, 2022). Didymocarpin, 5,8-dihydroxy-7-methoxyflavone, isopedicin, pediflavone, didymocarpin A, pedicellin, pedicin, pedicinin and methyl pedicin are the major polyphenolic constituents present in leaves of *D. pedicellatus*. (Saklani *et al.*, 2021) These chemical constituents vary among different populations depending not only upon the genotype but also influenced by the habitat conditions, altitudes and prevailing climatic conditions such as temperature, atmospheric pressure and humidity etc. It is necessary to see the impact of elevation on the polyphenolic content so as to identify the optimum yielding population that

can supply quality material for herbal industry. Further, the higher yielding population can also be used as elite planting material for cultivation purpose. Elevation also impacts the photosynthetic machinery, particularly chloroplast and chlorophyll pigment, which in turn affects the growth and other major metabolic processes besides affecting nutrient uptake and its assimilation (Mooney and Billings, 1961).

The current study aimed to investigate the effect of altitude on polyphenolic content, chlorophyll levels, protein content and concentrations of key nutrients (NPK) in plant samples collected from different altitudes in Uttarakhand. The study examined altitude-responsive traits of the plant to establish theoretical framework for comprehending its growth and adaptive mechanism to different environmental conditions aiming for sustainable agriculture.

MATERIALS AND METHODS

Collection of plant material and extract preparation

Plant samples of *D. pedicellatus* were collected from four different sites in Uttarakhand. Details of the elevation and GPS coordinates of the sites (A1, A2, A3 and A4) are given in Table 1. Voucher specimen was deposited in the herbarium of Department of Biological Sciences (GBPUH) with accession number 1441. Uprooted plant samples (along with the soil), were brought to the Plant Bioprospecting Laboratory in an aerated cool chamber (to keep the plant fresh). Leaves of the plants were immediately used for estimation of chlorophyll and protein content. Shade dried leaves were pulverized and methanolic extract was prepared using soxhlet extractor and stored at 4 °C for further use. All the chemicals used are of Hi Media make.

Estimation of total phenolics, flavonoids and tannins

Spectrophotometric estimation of total flavonoid content (TFC) was done using the method of Djeridane *et al* (2006) with Quercetin as the standard. Total phenolic content (TPC) was evaluated using Folin-Ciocalteu reagent (Wolfe *et al.*, 2003) with Gallic acid as the standard;

absorbance was taken at 665 nm and 765 nm respectively. The results were expressed in mg quercetin equivalent (QE) /g of extract for flavonoids and mg gallic acid equivalent (GAE) /g of extract for phenolic content. Total tannin content was estimated using protocol of Price and Butler (1977) with minor modifications. For this, diluted extracts were mixed with 0.1 M FeCl₃ and 8 mM potassium ferricyanide followed by incubation for 10 minutes at 27± 2°C. The optical density was measured at 720 nm using GENESYS 10S UV-VIS spectrophotometer (Thermo Scientific). Tannic acid standard curve was prepared for quantification of tannin content in the extract and results were expressed in mg tannic acid equivalent (TAE)/ g of extract.

Estimation of protein content

Spectrophotometric estimation of protein was done from the fresh leaves of *D. pedicellatus* following Bradford (1976) method using BSA (bovine serum albumin) for preparation of standard curve. The absorbance was recorded at 595 nm and protein content was expressed in mg BSA/g of fresh leaves.

Estimation of macronutrients

Leaf samples were used for evaluation of macronutrients (Nitrogen, Phosphorus and Potassium) using standard protocols (Okaebo *et al.*, 1993). Leaves were washed, shade dried and powdered for estimation of macronutrients. Nitrogen was quantified using Kjeldahl method (KJEL PLUS system, Pelican, India), by digestion, distillation and titration of leaf sample and content was expressed in mg /100 g dry weight. For phosphorus evaluation, a reaction mixture of ammonium molybdate, conc. H₂SO₄, and stannous chloride was added to the digested leaf samples. Absorbance was read at 420 nm using RAY LEIGH UV 2601 spectrophotometer. Absorbance vs. concentration standard curve of KH₂PO₄ was plotted and phosphorus content was estimated using the curve and expressed in mg/100 g dry weight. Potassium content was determined with digested sample using flame-photometer (128 Systronics). Potassium chloride (KCl) standard curve was prepared and concentration was read directly from the photometer.

Estimation of photosynthetic pigments

Spectrophotometric evaluation of chlorophyll a, b and carotenoids was done from the leaf samples of the plants collected from different altitudes using Hiscox and Israelstam (1979) adopted by Richardson *et al* (2002). Pre heated 1 ml dimethylsulfoxide (DMSO) per 10 mg of fresh leaf tissue was incubated for 30 min. at 65 °C. Absorbance was measured at 470, 645 and 663 nm, calibrating to zero with pure DMSO.

The following formula was used which was based on Arnon's (1949) equations, where:

$$\text{Chlorophyll } a = [(12.7 \times A_{663}) - (2.69 \times A_{645})] \times \frac{V}{1000 \times W}$$

$$\text{Chlorophyll } a = [(22.9 \times A_{645}) - (4.68 \times A_{663})] \times \frac{V}{1000 \times W}$$

$$\text{Carotenoids} = \left(\frac{1000 \times A_{470} \times 1.82 \text{ Chl } a + 85.02 \text{ Chl } b}{214} \right) \times \frac{V}{1000 \times W}$$

Where; V= volume of extract (ml), W= fresh weight of sample (g)

Statistical analysis

All the experiments were done in triplicates. Data was subjected to one way ANOVA and values were expressed in mean $\pm$ standard deviation using MS-Excel and Origin pro software. Tukey's mean comparison test was used to evaluate significance of data at $P < 0.05$.

RESULTS AND DISCUSSION

Total phenolics, flavonoids and tannin content

Total phenolics, flavonoid and tannin content analysed quantitatively across plant samples collected from different altitudes (A1 to A4 - where A1 is the lowest altitude, 700 m and A4 is the highest altitude, 1640 m), showed significant variations ($p < 0.05$) as depicted in Table 2. Total phenolic content ranged from 27.94 ± 0.35 mg GAE/g extract in A1 to 34.44 ± 0.43 mg GAE/g extract in A4 showing 23.26% increase with increasing altitude. The results revealed an increase in phenolic content with increasing altitude. Similar trend was observed in leaves of *Crataegus x sinaica*, *Thalictrum foliolosum*, *Tanacetum sinaicum* and *Cotoneaster orbicularis* that also showed elevated phenols with increasing elevation (Ibrahim *et al.*, 2022). *Asarum sieboldii* showed significant increase in total phenols

with increasing altitude (Pan *et al.*, 2023).

In contrast, total flavonoid content declined from 19.00 ± 1.33 mg QE/g extract (A1) to 2.88 ± 0.52 mg QE/g extract (A4) resulting in 84 % decrease with ascending elevation (Table 2). Similar trend was observed in *Polygonatum verticellatum* which also showed negative correlation with altitude ($r = -0.708$, $p < 0.01$) (Suyal *et al.*, 2019). Generally, flavonoid content increase with increasing altitude but this unusual trend suggests potential redistribution of resources towards other secondary metabolites during stress. Tannins also showed a marked increase of 63.75 % with increasing altitude, showing maximum tannin content at A4 (31.03 ± 0.85 mg TAE/g extract), suggesting an adaptive response to oxidative stress prevalent at higher elevations.

In the present study, phenols and tannin content showed an increasing trend with increasing altitude whereas flavonoid showed a reverse trend with increasing altitude. Rana *et al* (2020) showed increase in phenols, flavonoids and terpenoid content with increasing altitude in *Coleus forskohlii* collected from different altitudes of Garhwal region of Uttarakhand. Various abiotic stress like heavy metal, drought, high/low temperature and ultraviolet radiations elevate biosynthetic pathway of phenylpropanoid triggering increase of polyphenolic content in plants (Sharma *et al.*, 2016; Wang *et al.*, 2019; Nataraj *et al.*, 2022). Flavonoid pathway might be more dependent on other abiotic factors viz., UV light, topography of the habitat.

Total protein content

Protein also showed a significant upward trend from 27.06 ± 1.21 mg BSA /g (A1) to 74.61 ± 0.47 mg BSA /g FW (A4) giving an impressive 176 % increase in protein profile when moving up the hill, Table 2. Similar reported in *Kobresia pygmaea*, *Nepeta septemcrenata* and *Rosa arabica* yielding significantly high protein content at higher elevations (Li *et al.*, 2014; Hashim *et al.*, 2020). Dhatwalia *et al* (2024), also observed significant increase in protein content in *Rubus ellipticus* being minimum at 500 m and maximum at 2000 m. Protein accumulation actually fulfil the energy requirement of a plant in response to stress as well as scavenge

Table 1: Collection sites of *D. pedicellatus* in Uttarakhand, India

S.No.	Collection sites	Altitude (m)	Latitude	Longitude
1	Nainital (A1)	700	29° 17' 43.55" N	79° 32' 25.57" E
2	Nainital (A2)	1186	29° 20' 17" N	79° 29' 14" E
3	Chamoli (A3)	1370	30°15' 18 N	79°26' 48 E
4	Bageshwar (A4)	1640	29 ° 54' 49" N	79 ° 46' 34 " E

Where A1 is the lowest altitude and A4 is the highest altitude.

Table 2: Total phenolics, flavonoids, tannin and protein content in leaves of *D. pedicellatus* collected from different altitudes

Collection sites	TPC $\pm$ SD mg GAE/ g of extract	TFC $\pm$ SD mg QE/g of extract	TTC $\pm$ SD mg TAE/g of extract	Protein (mg BSA/g FW)
A1	27.94 $\pm$ 0.35 c	19.00 $\pm$ 1.33 a	18.95 $\pm$ 0.71 d	27.06 $\pm$ 1.21 d
A2	28.83 $\pm$ 0.79 c	10.26 $\pm$ 0.63 b	20.40 $\pm$ 1.80 c	50.44 $\pm$ 1.57 c
A3	31.06 $\pm$ 0.43 b	6.77 $\pm$ 0.56 c	25.55 $\pm$ 0.76 b	58.26 $\pm$ 0.86 b
A4	34.44 $\pm$ 0.43 a	2.88 $\pm$ 0.52 d	31.03 $\pm$ 0.85 a	74.61 $\pm$ 0.47 a
CV	0.017	0.085	0.047	0.021

Data are represented as mean $\pm$ standard deviation. Tukey's mean comparison at significance level $P < 0.05$. CV – Coefficient of variation. A1- 700 m, A2- 1186 m, A3- 1370 m and A4- 1640 m.

Table 3. Photosynthetic pigments in leaves of *D. pedicellatus* collected from different altitudes

Collection sites	Chlorophyll a (mg/ 100 g fresh weight)	Chlorophyll b (mg/ 100 g fresh weight)	Carotenoids (mg/100 g fresh weight)
A1	78.20 $\pm$ 0.95 a	35.12 $\pm$ 0.34 a	32.89 $\pm$ 0.70 a
A2	75.84 $\pm$ 0.65 b	31.95 $\pm$ 0.75 b	29.55 $\pm$ 0.56 b
A3	69.48 $\pm$ 0.55 c	24.97 $\pm$ 1.61 c	24.83 $\pm$ 0.89 c
A4	49.69 $\pm$ 0.44 d	21.08 $\pm$ 0.67 d	20.97 $\pm$ 0.17 d
CV	0.01	0.034	0.02

Data are represented as mean $\pm$ standard deviation. Tukey's mean comparison at significance level $P < 0.05$. CV – Coefficient of variation. A1- 700 m, A2- 1186 m, A3- 1370 m and A4- 1640 m.

Table 4: Nitrogen (N), phosphorus (P) and potassium (K) level in leaves of *D. pedicellatus* collected from different altitudes

Collection sites	Nitrogen (mg / 100 g)	Phosphorus(mg / 100 g)	Potassium(mg / 100 g)
A1	789.00 $\pm$ 1.00 a	214.00 $\pm$ 5.29 a	1315.00 $\pm$ 13 a
A2	624.67 $\pm$ 4.73 b	204.67 $\pm$ 6.43 a	989.67 $\pm$ 10.02 b
A3	484.33 $\pm$ 7.37 c	186.33 $\pm$ 5.51 b	633.67 $\pm$ 11.02 c
A4	351.67 $\pm$ 3.79 d	169.67 $\pm$ 9.07 b	556.67 $\pm$ 7.02 d
CV	0.009	0.035	0.012

Data are represented as mean $\pm$ standard deviation. Tukey's mean comparison at significance level $P < 0.05$. CV – Coefficient of variation. A1- 700 m, A2- 1186 m, A3- 1370 m and A4- 1640 m.

ROS as a defence response (Khan *et al.*, 2016).

Macronutrient content

Nutritional content of the leaves also significantly varied with the altitude (Table 4). Analysis of macronutrients in the leaves of *D. pedicellatus* collected from different altitudes showed significant decreasing trend of nitrogen (A1- 789.00 $\pm$ 1.00 to A4 - 351.67 $\pm$ 3.79) and potassium (A1- 1315.00 $\pm$ 13 mg /100 g, A4 – 556.67 $\pm$ 7.02 mg/100 g) with

rising elevation. Decrease in phosphorus, however (A1- 214.00 $\pm$ 5.29 mg/ 100 g, A4 - 169.67 $\pm$ 9.07 mg/100 g) was not very significant in relation to varying elevations. More than half of the nitrogen (55.43 %) and potassium content (57.67 %) decreased with increase of about 940 m altitude whereas phosphorous had a decrease of only 20.71 %. The results are in conformity with other recent studies on altitudinal variation in leaf nutrient content (Tsombou *et al.*, 2025). Chandra and Lata

(2022) also found decrease in potassium and phosphorous content in *Aconitum balfourii* and *Podophyllum hexandrum* with an exceptional increase in nitrogen content. In the present study, nitrogen decrease may be possibly due to over accumulation of protein content in the leaves of *D. pedicellatus* at higher altitudes. Phosphorus and potassium being involved in maintaining other metabolic processes for sustaining life at higher altitudes, also showed a decline. This decrease can be explained owing to comparatively lower soil temperature and hence decreased soil microbial activity causing poor nutrient absorption (Zhu *et al.*, 2024).

Photosynthetic pigments

Altitude and environmental conditions greatly impact photosynthesis and its pigments. The results revealed a significant decrease in chlorophyll a, b and carotenoids content with increasing altitude (36.46%, 39.98 % and 36.24 % respectively). Maximum chlorophyll-a content was 78.20 ± 0.95 at A1 elevation whereas lowest was 49.69 ± 0.44 at A4 elevation as shown in Table 3. Similarly, Chlorophyll b and carotenoids were maximum at A1 (35.12 ± 0.34 , 32.89 ± 0.70) and minimum at A4 (21.08 ± 0.67 , 20.97 ± 0.17) showing significant difference at $p < 0.05$ (Table 3). This trend aligns with the studies of Ahmad *et al.* (2018), Cui *et al.* (2018), Zhao *et al.* (2020) and Chandra and Lata (2022) and indicating decrease in all the photosynthetic pigments with increasing altitude which ultimately affect the photosynthetic capacity of the plants growing at higher altitude leading to slower growth and poor development. Low-temperature stress at higher elevations affect synthesis of chlorophyll pigments and also its ability to harvest light due to increased photo-oxidation (Ibrahim *et al.*, 2022).

CONCLUSION

This study reveals notable effect of altitude on various biochemical and physiological parameters (phenols, flavonoids, tannins, chlorophylls, macronutrients and proteins) of a plant. The results divulged that phenol, tannin and protein content of

the plants are directly proportional to altitude whereas photosynthetic pigments (chl a, chl b and carotenoids), flavonoids and macronutrients (nitrogen, phosphorus and potassium) decrease in content with elevation. Our findings reveal distinct variation patterns across four different altitudes, suggesting adaptive response of *D. pedicellatus* to changing environmental conditions of temperature, precipitation, humidity, soil microbial activity and soil nutrient profile. This research fills a gap by providing comprehensive data on how altitudinal variation influence not just nutrient status but the overall physiological profile of plants. The study will also be helpful in selecting the elite population and its adaptability to plan its sustainable propagation in suitable mountainous ecosystems.

ACKNOWLEDGMENTS

The authors would like to express their sincere thanks to Dean, CBSH and Director Research, G. B. Pant University of Agriculture & Technology, Pantnagar for providing necessary facilities to conduct this work. Council of Scientific and Industrial Research (CSIR), New Delhi is also thankfully acknowledged for the financial support to carry out the work.

REFERENCES

- Ahmad, K. S., Hameed, M., Hamid, A., Nawaz, F., Kiani, B. H., Ahmad, M. S. A., Deng, J., Ahmad, F., Hussain, I., and Fatima, S. (2018). Beating cold by being tough: Impact of elevation on leaf characteristics in *Phleum himalaicum* Mez., endemic to the Himalayas. *Acta Physiologiae Plantarum*, 40: 1–17.
- Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiology*, 24:1–15
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1–2):248–254.
- Chandra, S. and Lata, H. (2022). Effect of altitude

- on biochemical and photosynthetic characteristics of *Aconitum balfourii* and *Podophyllum hexandrum*: high value endangered medicinal herbs from Himalayas. *International Journal of Tropical Agriculture*, 40:195-203
- Cui, G., Li, B., He, W., Yin, X., Liu, S., Lian, L., Zhang, Y., Liang, W. and Zhang, P. (2018). Physiological analysis of the effect of altitudinal gradients on *Leymus secalinus* on the Qinghai-Tibetan Plateau. *PLoS ONE*, 13(11), e0202881.
- Dhatwalia, J., Kumari, A., Guleria, I., Kumar Shukla, R., Saleh, N. I., El-Nashar, H. A. and El-Shazly, M. (2024). Effect of altitude and harvest year on nutraceutical characteristics of *Rubus ellipticus* fruits. *Frontiers in Nutrition*, 11: 1419862.
- Djeridane, A., Yousfi, M., Nadjemi, B., Boutassouna, D., Stocker, P. and Vidal, N. (2006). Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chemistry*, 97:654–660.
- Dongdong, C., Qi, L., Zhe, L., Fuquan, H., Xin, C., Shixiao, X., Xinquan, Z. and Liang, Z. (2020). Variations of forage yield and nutrients with altitude gradients and their influencing factors in alpine meadow of Sanjiangyuan, China. *Journal of Soil Science and Plant Nutrition*, 20: 2164-2174.
- Du, B., Ji, H., Peng, C., Liu, X., and Liu, C. (2017). Altitudinal patterns of leaf stoichiometry and nutrient resorption in *Quercus variabilis* in the Baotianman Mountains, China. *Plant and Soil*, 413: 193–202.
- Goyal, P.K., Verma, S.K., Sharma, I., Sharma, A. and Sharma, A.K. (2015). Ethnobotanical and phytopharmacological updates on *Didymocarpus pedicellata*. *Indo American Journal of Pharmacological Science*, 5(08): 2747-2755.
- Hashim, A. M., Alharbi, B. M., Abdulmajeed, A. M., Elkelish, A., Hozzein, W. N. and Hassan, H. M. (2020). Oxidative stress responses of some endemic plants to high altitudes by intensifying antioxidants and secondary metabolites content. *Plants*, 9(7): 869.
- Hiscox, J.D. and Israelstam, G.F. (1979). A method for the extraction of chlorophyll from leaf tissue without maceration. *Canadian Journal of Botany*, 57(12): 1332-1334.
- Ibrahim, I. A., Jabbour, A. A., Abdulmajeed, A. M., Elhady, M. E., Almaroai, Y. A., and Hashim, A. M. (2022). Adaptive responses of four medicinal plants to high altitude oxidative stresses through the regulation of antioxidants and secondary metabolites. *Agronomy*, 12(12):3032.
- Khan, M. N., Mobin, M., Abbas, Z. K., and AlMutairi, K. A. (2016). Impact of varying elevations on growth and activities of antioxidant enzymes of some medicinal plants of Saudi Arabia. *Acta Ecologica Sinica*, 36(3): 141–148.
- Körner, C. (2007). The use of ‘altitude’ in ecological research. *Trends in Ecology & Evolution*, 22(11): 569–574.
- Li, X., Yang, Y., Ma, L., Sun, X., Yang, S., Kong, X., Hu, X. and Yang, Y. (2014). Comparative proteomics analyses of *Kobresia pygmaea* adaptation to environment along an elevational gradient on the central Tibetan Plateau. *PLoS ONE*, 9(6): e98410.
- Mooney, H.A. and Billings, W.D. (1961) Comparative physiological ecology of arctic and alpine population of *Oxyria digyna*. *Ecological Monographs*, 31: 1-29.
- Nanjala, C., Odago, W.O., Rono, P.C., Waswa, E.N., Mutinda, E.S., Oulo, M.A., Muema, F.W., Wanga, V.O., Mkala, E.M., Kuja, J. and Njire, M.M. (2022). A review on ethnobotany, phytochemistry, and pharmacology of the genus *Didymocarpus* wall. (Gesneriaceae). *Journal of Ethnopharmacology*. 295: 115404.
- Nataraj, N., Hussain, M., Ibrahim, M., Hausmann, A. E., Rao, S., Kaur, S., Khazir, J., Mir, B. A., and Olsson, S. B. (2022). Effect of altitude on volatile organic and phenolic compounds of *Artemisia brevifolia* Wall ex Dc. from the Western Himalayas. *Frontiers in Ecology and Evolution*, 10: 864728.
- Okaebo, J.R., Gathua, K.W. and Whoomer, P.L.

- (1993). *Laboratory method of soil and plant analysis: A working manual*. Nairobi, Kenya: Tropical Soil Biology and Fertility (TSBF) Programme, Pp. 22–29
- Pan, L., Yang, N., Sui, Y., Li, Y., Zhao, W., Zhang, L., Mu, L. and Tang, Z. (2023). Altitudinal variation on metabolites, elements, and antioxidant activities of medicinal plant asarum. *Metabolites*, 13(12): 1193.
- Price, M.L. and Butler, L.G. (1977). Rapid visual estimation and spectrophotometric determination of tannin content of sorghum grain. *Journal of Agricultural and Food Chemistry*, 25(6):1268–1273.
- Rana, P.S., Saklani, P. and Chandel, C. (2020). Research article influence of altitude on secondary metabolites and antioxidant activity of *Coleus forskohlii* root extracts. *Res. J. Med. Plants*, 14: 43–52.
- Richardson, A.D., Duigan, S.P. and Berlyn, G.P. (2002). An evaluation of noninvasive methods to estimate foliar chlorophyll content. *New phytologist*, 153(1):185–194.
- Saklani, A., Singh, D.C., Semwal, D. and Arya, R. (2021). *Didymocarpus pedicellata* R BR. A valuable herb for renal stones. *European Journal of Biomedical and Pharmaceutical Sciences*, 8(7): 268–272.
- Sharma, A., Thakur, S., Kumar, V., Kanwar, M. K., Kesavan, A. K., Thukral, A. K., Bhardwaj, R., Alam, P., and Ahmad, P. (2016). Pre-sowing seed treatment with 24-epibrassinolide ameliorates pesticide stress in *Brassica juncea* L. through the modulation of stress markers. *Frontiers in Plant Science*, 7:1569.
- Suyal, R., Rawat, S., Rawal, R. S., and Bhatt, I. D. (2019). Variability in morphology, phytochemicals, and antioxidants in *Polygonatum verticillatum* (L.) All. populations under different altitudes and habitat conditions in the Western Himalaya, India. *Environmental Monitoring and Assessment*, 191: 1–18.
- Tsombou, F.M., Al Dhanhani, A.S.S., Alhmoudi, A.M.S.A., Al Hmoudi, M.A.S.M., Ridouane, F.L., Alabdouli, M.F.A., Alhammadi, H.J., Alsamahi, M.E.A.A. and Mirza, S.B. (2025). Altitudinal influence on survival mechanisms, nutritional composition, and antimicrobial activity of *Moringa Peregrina* in the summer climate of Fujairah, UAE. *Scientific Reports*, 15(1): 5635.
- Wang, L. I., Shan, T., Xie, B., Ling, C., Shao, S., Jin, P. and Zheng, Y. (2019). Glycine betaine reduces chilling injury in peach fruit by enhancing phenolic and sugar metabolisms. *Food Chemistry*, 272:530–538.
- Wolfe, K., Wu, X. and Liu, R.H., (2003). Antioxidant activity of apple peels. *Journal of Agricultural and Food Chemistry*, 51: 609–614.
- Zhao, Y., Han, Q., Ding, C., Huang, Y., Liao, J., Chen, T., Feng, S., Zhou, L., Zhang, Z., Chen, Y. and Yuan, S. (2020). Effect of low temperature on chlorophyll biosynthesis and chloroplast biogenesis of rice seedlings during greening. *International Journal of Molecular Sciences*, 21(4):1390.
- Zhu, D., Zhou, Y., Peng, S., Hua, W., Luo, B. and Hui, D. (2024). Impacts of altitude on plant green leaf, fresh litter, and soil stoichiometry in subtropical forests. *Frontiers in Forests and Global Change*, 7: 1–12.

Received: April 16, 2025

Accepted: April 25, 2025