

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 23(1) January-April 2025]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology, Pantnagar



ADVISORY BOARD

Patron

Prof. Manmohan Singh Chauhan, Ph.D., Vice-Cancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. A.S. Nain, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. A.H. Ahmad, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.P.S. Gangwar, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Subhash Chandra, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Anil Kumar Gaur, Ph.D., Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A

Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Indu S Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

M.P. Pandey, Ph.D., Ex. Vice Chancellor, BAU, Ranchi & IGKV, Raipur, Director General, IAT, Allahabad, India

Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agri. & Tech., Pantnagar, India

Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

Editor-in-Chief

Manoranjan Dutta, Ph.D., Ex Head, Germplasm Evaluation Division, National Bureau of Plant Genetic Resources, New Delhi, India

Managing Editor

S.N. Tiwari, Ph.D., Professor (Retd.) & Ex-Director Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head

Brijesh Dumka, Developer & Programmer

CONTENTS

<i>In-silico</i> analysis of curcumin conjugates targeting the Wnt signaling pathway in Breast Cancer Stem Cells	1
KANCHAN GAIROLA and SHIV KUMAR DUBEY	
Investigating <i>in vitro</i> direct antagonistic effect of endophytic bacteria against <i>Alternaria brassicicola</i>	8
SHIVANGI KRISHNATRA and A. K SHARMA	
Impact of altitude on photosynthetic and biochemical profile of <i>Didymocarpus pedicellatus</i> R.Br.: an antiurolithiatic Himalayan herb	18
DIVYA and PREETI CHATURVEDI	
Impact of Integrated Nutrient Management (INM) on growth, yield, quality and soil fertility status in sugarcane-ratoon system	25
JYOTI PAWAR and DHEER SINGH	
Mapping and evaluation of soil macronutrient and micronutrient status in Muzaffarnagar district of India	32
RAUSHAN KUMAR and G. R. SINGH	
Study of shift in cropping pattern in northern dry zone of Karnataka	45
ASHWINI HEBBAR and SUMA A. P.	
Changing weather conditions during summer and early monsoon season in the <i>Tarai</i> region of Uttarakhand	51
SHIVANI KOTHIYAL and R.K. SINGH	
Nutrients enhancing flowering characteristics in Mango (<i>Mangifera indica</i> cv. Dashehari) under medium density planting	57
KULDEEP, ASHOK KUMAR SINGH and SHAILESH CHANDRA SHANKHDHAR	
Nutrients and antioxidants potential of star fruit (<i>Averrhoa carambola</i> L.)	66
ABHIMA K. MOORTHY and LAKSHMY P. S.	
Physico-chemical and anti-nutritional properties of predigested composite flour mix from corn and green gram	76
MANISHA RANI and ANJU KUMARI	
Standardisation and quality evaluation of coconut milk yoghurt	84
RINIYA THAJ and LAKSHMY P. S.	
Study on growth performance and morphometric traits of Chaugarkha goat kids in Almora hills of Uttarakhand	93
UMA NAULIA and B. N. SHAHI	

Bacterial isolates from tracheo-bronchial aspirates of healthy and pneumonic cattle ASMITA NARANG, CHARANJIT SINGH, MUDIT CHANDRA and DHIRAJ KUMAR GUPTA	99
Successful management of notoedric mange in two domestic cats: A case report ASMITA NARANG, GURPREET SINGH PREET, JASNIT SINGH and HARKIRAT SINGH	106
Dietary supplementation of formulated fish-specific mineral mixtures improved the growth, nutrient composition and health status of <i>Cyprinus carpio</i> fingerlings ABHED PANDEY, UDEYBIR SINGH CHAHAL and ANJU VIJAYAN	110
Impact of deep cryogenic treatment on microstructural and electrical properties of recycled aluminium alloys BIRENDRA SINGH KARKI and ANADI MISRA	118
Assessing farmers' attitudes and factors influencing livelihood diversification in Nainital District of Uttarakhand NEHA PANDEY, AMARDEEP and V.L.V. KAMESWARI	126
Perceived Benefits of Tribal Sub Plan (TSP) Project on tribal beneficiaries in Udham Singh Nagar District of Uttarakhand ARPITA SHARMA KANDPAL, JITENDRA KWATRA, VLV KAMESWARI and AMARDEEP	133

Physico-chemical and anti-nutritional properties of predigested composite flour mix from corn and green gram

MANISHA RANI¹ and ANJU KUMARI^{1*}

¹Centre of Food Science and Technology, Chaudhary Charan Singh Haryana Agricultural University, Hisar-125004 (Haryana)

*Corresponding author's email id: anjugaina@hau.ac.in

ABSTRACT: Mung bean (*Vigna radiata*), commonly called green gram, is a nutrient-rich legume from the Fabaceae family, widely consumed for its health benefits and versatility in savoury and sweet dishes. It is a valuable source of plant-based protein, dietary fibre, essential vitamins, and minerals, making it an integral part of balanced diets globally. Corn (*Zea mays*), a gluten-free cereal staple, is equally celebrated for its high content of carbohydrates, protein, and essential micronutrients such as magnesium, phosphorus, and vitamin B complexes. Despite their inherent nutritive properties, mung bean and corn exhibit anti-nutritional factors such as phytates and tannins that can reduce nutrient absorption and bioavailability. Enhancing these grains' nutritional profiles while addressing these limitations is crucial for their effective incorporation into health-focused diets. This study utilized germination and lactic acid fermentation as innovative processing techniques to improve the digestibility and bioavailability of nutrients in mung bean and corn. These processes have been shown to reduce anti-nutritional compounds, enrich grains with beneficial bioactive compounds, and enhance their overall nutritional quality. Nine composite flours were developed by blending different ratios of germinated and fermented flours of mung bean and corn. The physico-chemical properties, such as moisture content, nutrient composition, and texture, were analyzed alongside anti-nutritional factors to evaluate the suitability of these flours. The findings underline the significant potential of predigested flours for creating value-added recipes that cater to modern dietary needs, including high-protein, high-fiber, and gluten-free food products. These flours could be pivotal in addressing nutritional deficiencies, improving gut health, and developing functional foods to support better health outcomes.

Keywords: Composite flour, germination, fermentation, corn, mungbean

Composite flour is a blend of flours produced from cereals, roots, tubers, and legumes, with or without wheat flour. It promotes using indigenous agricultural products such as flour, and accepting these native crops would increase their usage via processing and value addition (Bello *et al.*, 2022). Eating whole grain cereals, legumes, and their products, including biscuits, prevents nutrition-related non-communicable diseases (Irondi *et al.*, 2024).

Green gram (*Vigna radiata*) is a grain legume that matures in 70–90 days and is also known as mungbean or golden gram in some places worldwide. It is commonly cultivated in the tropics and subtropics. It is consumed as cooked beans, broth, or pancakes and is an essential dietary source for people. Green gram seeds are high in protein (20.97–31.32%), and their carbohydrates are easier to digest, resulting in less flatulence in humans than other legumes (Nair *et al.*, 2013). It is a healthy legume that is widely consumed in South East Asia. It is

well-known for having a high digestibility and not causing flatulence, which is frequently associated with many legumes. Legumes have dietary fibre content ranging from 8 to 27.5% and soluble fibre content ranging from 3.3% to 13.8% (Eashwarage, 2017).

Cereals are an excellent source of vitamins and minerals, including fat-soluble vitamin E, an essential antioxidant. The cereal grains are an easy protein source as required by Recommended Daily Allowance (RDA) (Khatkar, 2005). Across the world, nutritional meals are produced from grains like wheat, rice, maize, oats, barley, rye, sorghum, millet, buckwheat, and quinoa, and these are then eaten whole, polished, or ground (Ding and Feng, 2025). Maize is one of the most widely cultivated plants in the world and the third most important cereal after rice and wheat (Gwirtz and Garcia-Casal, 2014). In addition, corn is a very interesting and nutritionally edible plant due to its richness in protein and certain minerals, carbohydrates and vitamins

(Shah *et al.*, 2011). However, anti-nutritional compounds can affect the digestibility of proteins and other nutrients (Gnanwa *et al.*, 2021). Corn flour develops and processes a vast range of foods and goods. “Food professionals have investigated the potential of various food ingredients/crops as well as appropriate technological processes to develop gluten-free products, but the quality of biscuits from blends of maize and soybean flour has not been assessed” (Okereke, 2023).

Germination is a simple, inexpensive, and environmentally friendly method of producing plant foods with functional properties. It enhances the nutritional and medicinal properties of plant foods by reducing antinutrients and increasing the accumulation of antioxidants, flavonoids, phenolic acids, and vitamins, thereby increasing the value of grains (Durovic *et al.*, 2021). The best protein sources are legumes, starch, vitamins, and minerals. They serve as meat substitutes or alternatives that give the vegetarian diet all the nutrients it needs due to their excellent nutritional value (Chaturvedi and Chakraborty, 2021). A final product with enhanced nutritional content, safety, stability, and sensory qualities may be produced through carefully regulated grain sprouting and fermentation processes (Wu and Xu, 2019).

Fermentation is one of the earliest methods used to preserve food. Native microbiota, such as moulds, aerobic spore formers, lactic acid bacteria (LAB), and enterobacteria, compete for nutritional deposits in cereal and pseudocereal grains. Many species of *Lactobacillus* are well-nourished by cereals and pseudocereals (Wronkowska *et al.*, 2023). Fermentation can convert nutrients such as carbohydrates and proteins into usable end products and increase the contents of essential amino acids, proteins, and minerals in foods. *Lactobacillus* fermentation has been reported to be one of the most cost-effective methods to maintain or improve food products' nutrition, organoleptic, safety, and shelf-life (Kwaw *et al.*, 2018). It can lead to partial or complete degradation of anti-nutritional factors, thereby improving protein quality and bioavailability of minerals (Gobbetti *et al.*, 2019). Although lactic acid bacteria (LAB) have been widely used in the food fermentation industry, studies on the effect

of *Lactobacillus* fermentation on the nutritional value of potato flour have not been reported. Since most LAB cannot directly degrade native starch, fermentation of starch-based foods is inefficient. The current study aimed to standardize the procedure for preparing a predigested mixture of green gram and corn and evaluating its quality.

MATERIALS AND METHODS

Pure culture of *Lactobacillus acidophilus* ATCC was procured from the Institute of Microbial Technology, Chandigarh, India or ATCC. The culture was maintained on slants and sub-cultured after every 30 days. Corn and green gram were procured from the Genetics and Plant Breeding Department, CCS Haryana Agricultural University, Hisar. We purchased other ingredients from the nearby market.

Treatments

- I. Germination: Grains were hygienist separately with 0.1% sodium hypochlorite (Grains: NaOCl ratio, 1:5 w/v) for 30 minutes and rinsed with distilled water. They were then soaked separately in distilled water (1:5 w/v grain to distilled water ratio) at room temperature and left overnight—hydrated grains germinated in the dark. The maximum time of germination was fixed by achieving 95% sprout grains.
- II. Fermentation: After cleaning, the grains were left to soak for 12 hours at room temperature in distilled water. The soaked grains were drained and dispersed in distilled water to make a slurry. *Lactobacillus acidophilus* (10^7 cells/ml) subjected the slurry to lactic acid fermentation. The germinated and fermented grains were dried at $55 \pm 5^\circ\text{C}$ and kept in airtight containers.
- III. Preparation of composite flour: Refined wheat flour (WF), mungbean flour (MF), germinated and fermented mungbean flour (GMF and FMF), corn flour, and germinated and fermented corn flour (GCF and FCF) were used to prepare composite flour. Nine different kinds of composite flour were made. One was prepared from whole wheat flour (WF) and three composites flours prepared from WF: MF: CF of 37.5:37.5:25 (C1), 25:25:50 of (C2) and 12.5:12.5:75 (C3), three germinated composite

flour consists WF:GMF: GCF of 37.5:37.5:25 (G4), 25:25:50 of (G5) and 12.5:12.5:75 (G6) and fermented composite flour consists WF: FMF: FCF of 37.5:37.5:25 (F7), 25:25:50 of (F8) and 12.5:12.5:75 (F9). The composite flours were passed through a 60 mesh size sieve for uniform mixing.

Proximate composition

Samples were evaluated for moisture content using a hot air oven, protein using the micro-Kjeldhal method with the KELPLUS Nitrogen estimation system, fat using the automatic SOCS plus Solvent extraction system, crude fibre using the automatic Fibre-Plus apparatus, and crude ash content using the standard AOAC method (2005). The Hulme and Narain(1931) methods calculated the total and reducing sugars. The sample's available iron was extracted (Rao *et al.*, 1978), and available zinc was extracted (Kim and Zemel, 1986). The amount of calcium was estimated using the AOAC (2005) standard procedure. Phytic acid was measured using the Haughand Lantzsch (1983) technique. Total polyphenols were extracted with the help of Singh and Jambunathan (1981).

The carbohydrate content was determined using the difference technique AOAC (2005) on a dry basis using the following formula: Total carbohydrates = 100 – (crude fat + crude protein + ash + crude fibre) Using the following formula, the energy was determined using the factorial technique by multiplying the sample's protein, carbohydrate, and fat contents by 4, 4, and 9, respectively.

Energy (K cal/100g) = 4.0 x protein (%) + 4.0 x carbohydrate (%) + 9.0 x fat (%).

Sensory evaluation- Chapattis were prepared from composite flours and evaluated for sensory parameters using a 9-point hedonic scale.

Statistical analysis: The data in this study were analyzed using SAS software and analysis of variance (ANOVA) procedures, following a completely randomized design.

RESULTS AND DISCUSSION

The present study tried to develop composite flour from predigested grain (Mungbean and Corn). The

composite flours' functional, chemical, and nutritional characteristics were examined. Nine types of composite flours (Figure 1.) were prepared by mixing various ratios of refined wheat flour (RF), mungbean flour (MF), germinated mungbean flour (GMF), fermented mungbean flour (FMF), corn flour (CF), germinated corn flour (GCF), and fermented corn flour (FCF).

Nutritional composition of RF, MF, GMF, FMF, CF, GCF and CCF

Proximate composition

Control samples (refined wheat flour, MF and CF), germinated (GMF and GCF), and fermented flour (FMF and FCF) were analyzed for proximate composition (Table 1).

There was a significant difference ($P < 0.05$) in the proximate composition of fermented, germinated, and refined wheat flours. The moisture content of a food indicates dry matter in that food (Adedajiet *al.*, 2014). Crude protein of GCF (20.52%) and crude fibre of GMF (17.03%) were the highest among all (Table 1). Fibre is essential to the diet that regulates




Fig. 1: Composite flour prepared by predigested Mungbean and Corn

bowel movement and weight. The increase in flour's crude fibre and ash contents is supported by Shah *et al.* (2011) and Devi *et al.* (2015), respectively. After sprouting, mung beans have increased antioxidant activity and are a good source of fibre, protein, vitamins, minerals and potential anti-cancer properties (Yadav, 2024).

CF, GCE and FCF contained (7.79, 7.90 and 6.49) moisture content; (1.12, 0.67 and 1.83%) ash; (5.93, 8.46 and 6.51 %) crude protein; (4.69, 4.42 and 4.41%) crude fat; (3.00, 3.03 and 3.16%) crude fibre and (84.98, 85.25 and 84.63%) carbohydrates respectively (Table 4.1). Similar findings in fermented corn flour (11.69% moisture, 7.63% crude protein, 4% crude fat, 1.62% ash, and 86.74% carbohydrate, respectively) were reported (Rahmawati *et al.*, 2018). The results show that GMF contained maximum crude protein and ash, whereas GCF contained maximum crude fat and total carbohydrates. Xu *et al.* (2020) reported that the GMF, which had high amounts of protein, ash, total sugar, lipid and fibre, could mix with the RF to produce gluten-free products with enhancement of the texture and nutritional quality of the final product. Pulses are a great protein, fibre, vitamins, and minerals source. Two examples of pulses are green gram and gram. The effect of adding sprouted green gram and sprouted gram flour to biscuits made with refined flour. When sprouted flour is added, the biscuits' protein content rises, but their fat and carbohydrate contents fall (Ding and Feng, 2025). Disrupting these interactions during fermentation and germination releases nutrients and phytochemicals, making them available to digest

Table 2: Mineral content in different composite flours

Flour	Iron(mg/100g)	Zinc(mg/100g)	Calcium(mg/100g)
WF	1.40±0.03 ^c	1.39±0.01 ^c	51.66±23.62 ^{cd}
MF	2.15±0.07 ^b	2.20±0.02 ^{bc}	83.66±8.50 ^{ab}
GMF	1.95±0.08 ^c	1.86±0.05 ^d	89.00±15.52 ^a
FMF	2.40±0.07 ^a	2.40±0.14 ^a	91.33±3.21 ^a
CF	2.07±0.05 ^b	2.15±0.08 ^c	31.66±7.63 ^d
GCF	1.68±0.06 ^d	1.73±0.20 ^d	37.33±6.80 ^d
FCF	2.34±0.02 ^a	2.36±0.05 ^{ab}	61.66±17.38 ^{bc}
CD	0.1	0.18	23.76

Values are mean ± SE of three independent determinations.

enzymes (Nkhata *et al.*, 2018). As expected, the fermentation of legumes degrades carbohydrates while increasing the concentration of soluble carbohydrates more easily absorbed by the gut (Adebo *et al.*, 2022). The composition of mungbean and corn changed after germination and fermentation.

Acidity and pH

The pH was an essential factor influencing overall qualities like colour, flavour and texture (Starowicz




Fig. 2: Acidity and pH of flour

Table 1: Proximate composition of different germinated and fermented flour

Flour	Moisture (%)	Ash (%)	Crude protein (%)	Crude fat (%)	Crude fibre (%)	Total carbohydrates (%)	Energy (kcal/100g)
WF	12.55±1.78 ^a	1.26±0.78 ^c	13.23±3.97 ^{ab}	0.75±0.79 ^{bc}	11.83±0.20 ^a	72.91±3.75 ^a	323.23±26.65 ^a
MF	8.65±5.25 ^a	3.75±1.85 ^a	11.52±3.26 ^b	1.04±0.47 ^c	16.53±5.24 ^a	66.93±5.29 ^a	321.45±23.65 ^a
GMF	10.17±7.50 ^a	3.87±0.08 ^a	12.99±3.25 ^{ab}	1.01±0.40 ^c	17.03±5.74 ^a	65.08±2.78 ^a	358.37±51.85 ^a
FMF	8.43±4.28 ^a	3.27±1.19 ^{ab}	15.17±5.26 ^{ab}	1.09±1.02 ^c	8.50±12.05 ^a	71.95±6.48 ^a	356.61±64.67 ^a
CF	10.70±4.34 ^a	1.12±0.69 ^c	17.51±3.15 ^{ab}	3.52±1.58 ^{ab}	14.13±14.85 ^a	63.70±12.50 ^a	376.48±17.90 ^a
GCF	10.75±4.35 ^a	1.69±0.45 ^{bc}	20.52±8.91 ^a	4.92±3.97 ^a	10.33±0.11 ^a	62.51±12.81 ^a	366.16±6.98 ^a
FCF	7.90±2.68 ^b	1.83±0.17 ^{bc}	17.22±3.72 ^{ab}	4.08±1.75 ^a	11.73±0.37 ^a	65.12±2.41 ^a	351.38±5.35 ^a
CD	8.12	1.64	8.58	3.1	13.67	13.54	63.56

Values are mean ± of three independent determinations

WF = Refined wheat flour, MF= Mungbean flour, GMF= Germinated mungbean flour FMF= Fermented mungbean flour CF= Corn flour, GCF= Germinated corn flour, FCF = Fermented corn flour

and Zielinski, 2019). Figure 2 shows a significant ($P < 0.05$) difference in pH and acidity among the various flours.

FMF contained higher acidity than MF and FCF, but GMF contained higher pH than MF and FCF. FCF is more acidic than CF and GCF. Fermented cassava flour with *L. plantarum* strain (LP) had the highest total titratable acidity, while commercial cassava flour had the lowest (Isa *et al.*, 2021).

Fermented flour may have a pH drop as a result of the fermentation process producing different organic compounds such as lactic, acetic acid, and ethanol, or a rise in acidity due to an increase in the concentration of fatty acids, phosphoric acids, H^+ , and carboxy groups of protein amino acid (Offiah *et al.*, 2017).

Minerals

All composite flours were analyzed for minerals (Table 2). There was a significant difference ($P < 0.05$) found among the minerals (Fe, Zn, and Ca) observed in each flour. Similar results in the calcium content of mungbean were also studied (Ohanenye *et al.*, 2020).

Essential nutrients, including potassium, phosphorus, zinc, calcium, iron, thiamine, niacin, vitamin B6, and folate, are abundant in corn flour (Arshad *et al.*, 2023). FMF and FCF stand out with the highest iron content, surpassing 2.3 mg/100g, while FMF also ranks highest in zinc and calcium content. This suggests that fermentation may play a positive role in enhancing mineral bioavailability. Mungbean fermentation or germination increased the accessibility of in vitro iron, zinc, and calcium (Nkhata *et al.*, 2018). Calcium, iron, and zinc concentrations enhanced when grass peas were fermented with lactic acid (Adebo *et al.*, 2022). In the present experiment, FCF contained maximum iron, zinc and calcium (Table 2). During fermentation, dry matter is lost whenever bacteria break down protein and carbohydrates, which increases mineral concentration (Nkhata *et al.*, 2018). The fermentation process improved mineral availability due to the breakdown of oxalates and phytase.

Protein bioavailability, particularly that of thiamine, iron, and calcium, increases dramatically during

germination. Additionally, it helps to increase the in vitro digestibility of starch and protein in dehulled and germinated green grams, cowpeas, lentils, and chickpeas (Starowicz and Zielinski, 2019).

Sugars

Figure 3 shows a significant difference ($P < 0.05$) in total and non-reducing sugar throughout all flours. The results showed that FCF contained higher non-soluble sugar than other flour.

MF, GMF, and FMF contained 4.40, 11.58, and 4.67 g/100 g total soluble sugar and 2.71, 3.99, and 3.75 g/100g non-reducing sugar. Compared to MF, FMF, and GMF, GMF had a higher total soluble and non-reducing sugar content. Germination significantly increased the amount of sugar in green gram (Chandra *et al.*, 2015; Nkhata *et al.*, 2018; Yadav 2024).

In the present study, the total sugar of CF, GCF, and FCF was 12.05, 16.04, and 13.06 g/100g, respectively. The enzymatic breakdown of starch in simple sugars during germination results in a significant increase in both total and non-reducing sugar. When monosaccharides hydrolyzed starch, sugar concentration in pulses increased (Starowicz and Zielinski, 2019). Germination of mung beans increased total sugars by germination (Chandra *et al.*, 2015; Yadav 2024).

Anti-nutritional factors

Among composite flours, GMF contained higher phytic acid and polyphenols. Phytic acid and polyphenol content increased in germinated mungbean (Chandra *et al.*, 2015; Yadav 2024). For




Fig. 3: Total soluble sugar and non-reducing sugar in flour

Table 3: Phytic acid and polyphenol content of different composite flours

Flour	Phytic acid (mg/100 g)	Polyphenols (mg/100 g)
WF	273.33±103.09 ^{bcd}	369.33±129.26 ^a
MF	401.73±140.24 ^a	313.86±72.71 ^a
GMF	416.40±58.62 ^b	300.80±38.06 ^{ab}
FMF	363.73±127.64 ^{bc}	249.33±79.05 ^b
CF	152.86±8.35 ^d	55.65±13.67 ^c
GCF	253.73±22.54 ^{cd}	51.09±16.07 ^c
FCF	218.26±56.48 ^{cd}	52.10±17.11 ^c
CD	153.52	115.47

Values are mean ± SE of three independent determinations

all flours, there was a significant ($P<0.05$) difference between the anti-nutritional (phytic acid and polyphenols) (Table 3).

Results indicated that GMF contained higher phytic acid and polyphenols than fermented treatments. During germination, the total phenolic contents of barley and mung beans increased significantly (Chandra *et al.*, 2015; Yadav 2024; Gupta and Bhatt 2024). The process of fermentation results in a significant decrease in phytic acid, polyphenols, and other antinutrients because the fermented dough has a low pH, which is optimal for phytase activity. Lectins and other proteinaceous anti-nutritional agents are affected by microbial proteolysis. The disulfide bonds should be broken to cause structural disruption, which is necessary for proteolysis. Several microbes have shown glutathione reductase activity. During the fermentation of legumes, thiol exchange processes catalyzed by glutathione are necessary to break down lectins (Arshad *et al.*, 2023).

The nutritional value of foods based on cereals is increased by the fermentation of cereal flour, which breaks down anti-nutritional factors. Fermented cereals with reduced tannin and phytic acid concentrations have more bioavailable minerals, which boosts the food's nutritional value (Adebo *et al.*, 2022). Increases in total phenolics and flavonoids were found in fermented cereals due to higher enzyme activity, and there was a positive correlation between total phenolics and antioxidant activity was also observed in previous studies (Saharan *et al.*, 2017). A higher amount of phytate consumption leads to a lack of certain minerals, such as zinc, magnesium, calcium, copper, and manganese, it hurts the minerals' bioavailability (Nkhata *et al.*, 2018).

Sensory value enhanced due to reduced anti-nutritional factor after fermentation increases mineral availability.

CONCLUSION

Refined wheat flour remains a staple in many cereal-based recipes; however, enhancing its nutritional profile by incorporating germinated and fermented cereal and pulse flours presents a promising opportunity. The significant increase in protein, calcium, iron, and zinc observed in fermented mung bean and corn flours highlights their potential to address nutrient deficiencies and promote health benefits. Chapattis prepared using specific blended flours (C2, C3, G5, and F9) exhibited desirable sensory characteristics, demonstrating the feasibility of incorporating composite flours into daily diets. Looking ahead, the application of composite flours holds significant promise in the development of diverse value-added products, such as protein-enriched breads, gluten-free baked goods, and nutritionally fortified snacks. Future research could explore optimizing blending ratios, extending shelf life, and assessing consumer preferences further to enhance these innovative flours versatility and market potential. By leveraging these advancements, composite flours can address global nutrition challenges and foster functional foods tailored to modern health-focused lifestyles.

REFERENCES

- Adediji, O. E., Oyinloye, O. D., and Ocheme, O. B. (2014). Effects of germination time on the functional properties of maize flour and its cookies' gelatinization degree. *African Journal of Food Science*, Pp. 8, 42–47.
- Adebo, J. A., Njobeh, P. B., Gbashi, S., Oyediji, A. B., Ogundele, O. M., Oyeyinka, S. A., and Adebo, O. A. (2022). Fermentation of cereals and legumes: Impact on nutritional constituents and nutrient bioavailability. *Fermentation*, 8(2): 63.
- AOAC (2005). Official Methods of Analysis of the Association of Analytical Chemists International, 18th ed. Gaithersburg, MD

- USA Official methods.
- Arshad, N., Akhtar, S., Ismail, T., Saeed, W., Qamar, M., Özogul, F. and Rocha, J. M. (2023). The Comparative Effect of Lactic Acid Fermentation and Germination on the Levels of Neurotoxin, Antinutrients, and Nutritional Attributes of Sweet Blue Pea (*Lathyrus sativus* L.). *Foods*, 12(15): 2851.
- Bello, F. A., Oladeji, B. S., and Akpan, P. S. (2022). Effect of green gram flour addition on the chemical composition and pasting properties of cocoyam flour as potential ingredients in food product development. *Acta Scientiarum Polonorum Technologia Alimentaria*, 21(3): 329-336.
- Chandra, S., Kumar, S., Vaishali, P., and Kumari, D. (2015). Effect of incorporation of rice, potato and mung flour on the physical properties of composite flour biscuits. *South Asian Journal Food Technology Environment*, 1(1), 64–74.
- Chaturvedi, S., and Chakraborty, S. (2021). Review on potential non-dairy synbiotic beverages: A preliminary approach using legumes. *International Journal of Food Science and Technology*, 56(5): 2068-2077.
- Devi, C. B., Kushwaha, A., and Kumar, A. (2015). Sprouting characteristics and associated changes in nutritional composition of cowpea (*Vigna unguiculata*). *Journal of Food Science and Technology*, 52: 6821-6827.
- Ding, J. and Feng, H. (2025). Controlling germination process to enhance the nutritional value of sprouted grains. In *Sprouted Grains*. Woodhead Publishing, Pp. 335-358.
- Đurovic, V., Radovanovic, M., Mandic, L., Knežević, D., Zornic, V., and Đukic, D. (2021). Chemical and microbial evaluation of biscuits made from wheat flour substituted with wheat sprouts. *Food Science and Technology International*, 27(2): 172-183.
- Eashwarage, I. S. (2017). Dietary fibre, resistant starch, and eleven commonly consumed legumes are Horse Gram. *Research Journal of Chemical*, 7(2): 1–7.
- Gobbetti, M., De Angelis, M., Di Cagno, R., Calasso, M., Archetti, G., and Rizzello, C. G. (2019). Novel insights on the functional/nutritional features of the sourdough fermentation. *International Journal of Food Microbiology*, 302: 103-113.
- Gnanwa, M. J., Fagbohoun, J. B., Ya, K. C., Blei, S. H., and Kouame, L. P. (2021). Assessment of minerals, vitamins and functional properties of flours from germinated yellow maize (*Zea mays* L.) seeds from Daloa (Côte D'Ivoire). *Int J Food Sci Nutr Eng*, 11(2): 35-42.
- Gwirtz, J. A., and Garcia Casal, M. N. (2014). Processing maize flour and corn meal food products. *Annals of the New York Academy of Sciences*, 1312(1): 66-75.
- Haugh, W., and Lantzsch, H.J. (1983). Sensitive method for the rapid determination of phytate in cereals and cereal products. *Journal Science Food and Agriculture*, 34:1423-1426.
- Hulme, A. C., and Narain, R. (1931). The ferricyanide method for the determination of reducing sugars: A modification of the Hagedorn-Jensen-Hanest technique. *Biochemical Journal*, 25(4): 1051.
- Irondi, E. A., Olatoye, K. K., Abdulameed, H. T., Aliyu, O. M., Ajani, E. O., and Ogbebor, O. F. (2024). Physicochemical, in vitro starch digestibility and sensory characteristics of biofortified yellow maize-cowpea composite flours and biscuits. *Food Production, Processing and Nutrition*, 6(1): 15.
- Isa, N. L. M., Kormin, F., Iwansyah, A. C., Desnilasari, D., and Hesani, A. (2021, April). Physicochemical properties and characterization of fermented cassava flour by lactic acid bacteria. In *IOP Conference Series: Earth and Environmental Science* (Vol. 736, No. 1, p. 012023). IOP Publishing.
- Kim, H., and Zemel, M. B. (1986). In vitro estimation of the potential bioavailability of calcium from sea mustard (*Undaria pinnatifida*), milk, and spinach under simulated normal and reduced gastric acid conditions. *Journal*

- of Food Science, 51(4), 957-959.
- Khatkar BS (2005). Trends in cereal processing. *Bev. Food World*. 32(11):65-67.
- Kwaw, E., Ma, Y., Tchabo, W., Apaliya, M. T., Wu, M., Sackey, A. S., ... and Tahir, H. E. (2018). Effect of lactobacillus strains on phenolic profile, color attributes and antioxidant activities of lactic-acid-fermented mulberry juice. *Food chemistry*, 250, 148-154.
- Nair, R. M., Yang, R. Y., Easdown, W. J., Thavarajah, D., Thavarajah, P., Hughes, J. D., Keatinge, J. D. (2013). Biofortification of mungbean (*Vigna radiata*) as a whole food to enhance human health. *J. Sci. Food Agric.*, 93, 1805–1813.
- Nkhata, S. G., Ayua, E., Kamau, E. H., and Shingiro, J. B. (2018). Fermentation and germination improve the nutritional value of cereals and legumes through the activation of endogenous enzymes. *Food Science and Nutrition*, 6(8), 2446-2458.
- Offiah, V. O., Abu, J. O., and Yusufu, M. I. (2017). Effect of co-fermentation on the chemical composition and sensory properties of maize and soybean complementary flours. *International Journal Innov. Food Science Technology*, 1, 18-28.
- Ohanenye, I. C., Tsopmo, A., Ejike, C. E., and Udenigwe, C. C. (2020). Germination as a bioprocess for enhancing the quality and nutritional prospects of Legume proteins. *Trends in Food Science and Technology*, 101, 213-222.
- Okereke GO. Quality evaluation of modified starches from white yam, trifoliate yam, sweet potato and utilization in wheat *Moringa oleifera* seed-based bakery products [PhD thesis]. Makurdi, Nigeria: Benue State University; 2023.
- Rahmawati, R., Maulani, R. R., and Saputra, D. (2018). Chemical properties, particle shape, and size of fermented local white corn flour of animal fs variety. *Journal Technology*, 80(5):155-161.
- Rao, B. N., and Prabhavathi, T. (1978). An in vitro method for predicting the bioavailability of iron from foods. *The American Journal of Clinical Nutrition*, 31(1): 169–175.
- Singh and Jambunathan, R. (1981). Studies on desi and kabull chickpea (*Ciceret al.*) cultivars: levels of protease inhibitors, levels of polyphenolic compounds and in vitro protein digestibility. *Journal of Food Science*, 46(5), 1364-1367.
- Shah, S. A., Zeb, A., Masood, T., Noreen, N., Abbas, S. J., Samiullah, M., and Muhammad, A. (2011). Effects of sprouting time on biochemical and nutritional qualities of mungbean varieties. *Afr. J. Agric. Res.*, 6: 5091-5098.
- Saharan, P., Sadh, P. K., and Duhan, J. S. (2017). Comparative assessment of effect of fermentation on phenolics, flavanoids and free radical scavenging activity of commonly used cereals. *Biocatalysis and Agricultural Biotechnology*, 12: 236-240.
- Starowicz, M., and Zieliński, H. (2019). How does the Maillard reaction influence the sensorial properties (colour, flavour and texture) of food products? *Food Reviews International*, 35(8): 707-725.
- Wu, F., and Xu, X. (2019). Sprouted grains-based fermented products. In *Sprouted Grains*. AACC International Press, Pp143-173.
- Wronkowska, M., Wiczowski, W., Topolska, J., Szawara-Nowak, D., Piskuba, M. K., and Zieliński, H. (2023). Identification and Bioaccessibility of Maillard Reaction Products and Phenolic Compounds in Buckwheat Biscuits Formulated from Flour Fermented by *Rhizopus oligosporus* 2710. *Molecules*, 28(6): 2746.
- Xu, J., Zhang, Y., Wang, W., and Li, Y. (2020). Advanced properties of gluten-free cookies, cakes, and crackers: A review. *Trends in Food Science and Technology*, 103: 200-213.
- Yadav, P. (2024). The nutritional importance of germinated wheat flour, mung beans, and beetroot: A critical review.

Received: January 23, 2025

Accepted: April 19, 2025