

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 23(2) May-August 2025]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology, Pantnagar



ADVISORY BOARD

Patron

Prof. Manmohan Singh Chauhan, Ph.D., Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. A.S. Nain, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. A.H. Ahmad, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Lokesh Varshney, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Subhash Chandra, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Ramesh Chandra Srivastava, Ph.D., Dean, College of Basic Sciences & Humanities, G.B.P.U.A.T., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A

Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Indu S Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

M.P. Pandey, Ph.D., Ex. Vice Chancellor, BAU, Ranchi & IGKV, Raipur, Director General, IAT, Allahabad, India

Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agri. & Tech., Pantnagar, India

Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

Editor-in-Chief

Manoranjan Dutta, Ph.D., Ex Head, Germplasm Evaluation Division, National Bureau of Plant Genetic Resources, New Delhi, India

Managing Editor

S.N. Tiwari, Ph.D., Professor (Retd.) & Ex-Director Research

G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head

Brijesh Dumka, Developer & Programmer

PANTNAGAR JOURNAL OF RESEARCH

Vol. 23(2)

May-August, 2025

CONTENTS

Bioaccumulation of heavy metals in soils and <i>Telfairia occidentalis</i> leaf grown around a river bank and dump site	139
ORHUE, E. R., EMOMU, A., JUDAH-ODIA, S. A., AIGBOGHAEBHOLO, O. P. and NWAEKE, I. S.	
Evaluation of maize cultivars for spring season in Indo-Gangetic plain of India	149
AMIT BHATNAGAR, N. K. SINGH and R. P. SINGH	
Weed management approaches for improving maize productivity in <i>Tarai</i> Belt of India	157
AKHILESH JUYAL and VINEETA RATHORE	
Effect of <i>Aloe vera</i> based composite edible coatings in retaining the postharvest quality of litchi fruits (<i>Litchi chinensis</i> Sonn.) cv. Rose Scented	163
GOPAL MANI, OMVEER SINGH and RATNA RAI	
Effect of chemical treatments on seed yield and quality in parthenocarpic cucumber (<i>Cucumis sativus</i> L.)	178
DHIRENDRA SINGH and UDIT JOSHI	
Assessment of chrysanthemum (<i>Dendranthema grandiflora</i> Tzvelev) varieties for their suitability for flower production under <i>Tarai</i> region of Uttarakhand	183
PALLAVI BHARATI and AJIT KUMAR KAPOOR	
Population dynamics of brown planthopper and mirid bug in relation to weather factors in the <i>Tarai</i> region	194
DEEPIKA JEENGAR and AJAY KUMAR PANDEY	
Influence of weather parameters on the population dynamics of Papaya mealybugs, <i>Paracoccus marginatus</i> and its natural enemies in Pantnagar, Uttarakhand	200
DIPTI JOSHI and POONAM SRIVASTAVA	
<i>In vitro</i> phosphate solubilizing and phyto stimulating potential of Rhizospheric <i>Trichoderma</i> from Hilly areas of Kumaun Region	208
DIVYA PANT and LAKSHMI TEWARI	
Economics of interventions and diversifications in existing farming systems in hills of Uttarakhand	221
DINESH KUMAR SINGH, AJEET PRATAP SINGH and ROHITASHAV SINGH	
Brucellosis surveillance and reproductive performance in an organized dairy herd of Uttarakhand: A seven-year retrospective analysis (2018–2024)	227
ATUL YADAV, SHIVANGI MAURYA, MAANSI and AJAY KUMAR UPADHYAY	
Effects of nanosilver administration on immune responses in Wistar Rats	230
NEHA PANT, R. S. CHAUHAN and MUNISH BATRA	

Antibacterial activity of Clove bud extract on MDR bacteria KANISHK A. KAMBLE, B. V. BALLURKAR and M. K. PATIL	240
Effect of iron oxide and aluminium oxide nanoparticles on biochemical parameters in Wistar rats NISHA KOHLI and SEEMA AGARWAL	247
Comprehensive case report of a mast cell tumor in a dog: clinical, cytological and histopathological analysis SWASTI SHARMA, SONALI MISHRA and GAURAV JOSHI	257
Evaluation of <i>In vitro</i> digestibility, functional and sensory characteristics of pre-digested corn and mungbean composite flour MANISHA RANI and ANJU KUMARI	261
Prevalence and public health correlates of constipation among adults in U. S. Nagar, Uttarakhand AKANKSHA SINGH, RITA SINGH RAGHUVANSHI and APURVA	270
Formulation and quality assessment of cheeses enriched with sapota pulp DELGI JOSEPH C. and SHARON, C. L.	279
Application of RSM for optimizing 7-day fermentation conditions in rice wine production RIYA K ZACHARIA, ANEENA E. R and SEEJA THOMACHAN	289
Investigating the mechanical properties and water absorption behavior of hemp-based natural fiber-reinforced bio-composites for humidity-resistant applications DEEPA SINGH and NEERAJ BISHT	303
Evaluating the performance of a forced convection solar drying system for chhurpi: A comparative analysis with traditional drying techniques SYED NADEEM UDDIN, SANDEEP GM PRASAD and PRASHANT M. DSOUZA	317
Digitization of G. B. Pant University Herbarium (GBPUH) and development of Virtual Herbarium Pantnagar, Uttarakhand (INDIA) RUPALI SHARMA, DHARMENDRA SINGH RAWAT and SANGEETA JOSHI	326
Constraints grappled with by rural communities during the implementation of Viksit Krishi Sankalp Abhiyan 2025 in Udham Singh Nagar District ARPITA SHARMA KANDPAL, B. D. SINGH, AJAY PRABHAKAR, SWATI and MEENA AGNIHOTRI	332

Bioaccumulation of heavy metals in soils and *Telfairia occidentalis* leaf grown around a river bank and dump site

ORHUE, E. R., EMOMU*, A., JUDAH-ODIA, S. A., AIGBOGHAEBHOLO, O. P. and NWAEKE, I. S.

Department of Soil Science, Faculty of Agriculture, University of Benin, Nigeria

**Corresponding author's email id: adams.emomu@uniben.edu*

ABSTRACT: River bank and dumpsites activities are threats to water and soil quality, as elevated heavy metal levels in soils around river banks and dumpsites may result in heavy metal entry into the food chain. This study aimed at accessing the concentration of heavy metals in soils and *Telfairia occidentalis* leaf grown around Temboga river bank and Oto-fure dumpsite, in Benin City, Nigeria. Surface (0-15 cm) soils and *T. occidentalis* leaf samples were collected at equidistance from Temboga river bank and Oto-fure dumpsite in 3 replicates. The soil samples were analyzed for some physical, chemical and heavy metal properties while *T. occidentalis* leaves were analyzed for Ni, Pb, Co, Cr, Se and Cd content in soil science analytical laboratory, University of Benin. Result showed that soil heavy metals content varied with distance from the river bank and dumpsite, Ni and Cr has reached toxic levels, while Pb, Cd, Se and Co accumulation are at potential toxic level in *T. occidentalis* leaf, with Ni content of 31.50, 13.00 and 12.50 mg kg⁻¹, Cr content of 8.92, 4.36 and 1.77 mg kg⁻¹ at dumpsite, 60 m and 120 m away from dumpsite, respectively. It could be concluded that *T. occidentalis* leaf grown around the dumpsite was contaminated with Ni and Cr, while *T. occidentalis* leaf grown around river bank, contained elevated levels of Co, all these will pose health risk to food safety and humans consuming *T. occidentalis* leaves grown around the river bank and dumpsite studied.

Keywords: Buildup, contamination, fluted pumpkin, health risk, permissible levels, toxic

Heavy metal bioaccumulation is the process through which toxic metals and chemical compounds become bonded inside a cell structure (Nnabueze *et al.*, 2023). The accumulation can lead to significant toxic effects, posing health risks to the organism itself and other organisms higher in the food chain, through bio-magnification. Heavy metals have their way into the soil and water either through, natural process of rock weathering and or anthropogenic activities (Bhagwat, 2019; Edelstein and Ben-Hur, 2018; Islam *et al.*, 2019; Xia *et al.*, 2021). The use of fertilizers, pesticides, and irrigation with contaminated water can introduce heavy metals into agricultural soils (Amadi *et al.*, 2020). When injurious metals are introduced into soils at certain levels, it could affect soil chemical and biological (Friedlova, 2010; Wang *et al.*, 2024), and deteriorate soil fertility properties (Rahi *et al.*, 2022). Various metal transfer mechanism could lead to heavy metal uptake by crop plants (Skuzza *et al.*, 2022). This can result in entry into the food chain and pose health risks to humans and animals (Amadi *et al.*, 2020). Due to nonbiodegradability, high toxicity and accumulation

through food chain, heavy metal pollution can affect ecosystem functions, food security and human health risk (Acharya, 2024). Health risks, including kidney damage, neurological disorders, and cancers are often associated with consuming foods containing elevated levels of injurious metals (Akanke *et al.*, 2021). While River banks are lands that flank a river and serves as boundary between flowing water and terrestrial environment, a dumpsite is an area of land used to dispose, store and manage waste. These land uses can results in certain environmental and human hazards (Ayoola, 2021). Some farmers now grow crops like *Telfairia occidentalis* in soils around river banks and dumpsites without knowledge of the heavy metals content in the soils and crops grown. Although several studies have been conducted on heavy metals status of crops grown along river banks (Adefarati *et al.*, 2024; Toheeb *et al.*, 2024) and dumpsites (Lissy and Madhu, 2020) soils and their effects on humans and animals health, but only little studies have examine heavy metal status in soils and *T. occidentalis* leaves around river banks and dumpsites in Benin metropolis. Hence, this study

aimed at evaluating heavy metal (Ni, Pb, Co, Cr, Se and Cd) levels in soils and *T. occidentalis* leaf grown around a Temboga river bank and Oto-fure dumpsite in Benin City, Nigeria

MATERIALS AND METHODS

The current study was carried out in areas where vegetables are cultivated for commercial purpose in Benin metropolis, Edo state, Nigeria, to evaluate heavy metal content of soils and *Telfairia occidentalis* (Fluted pumpkin) leaf grown around Temboga river bank and Oto-fure dumpsite. Temboga river bank site lies between latitude 6°21'28"N and longitude 5°38'56"E, sloppy and swampy closest to the river and planted to various vegetables, including *T. occidentalis*, *Amaranthus hybridus* (green leaf) and *Ocimum gratissimum* (scent leaf), the area is also used for fish farming activities. The soil in the area has a history of being treated with fertilizers (organic, inorganic) and herbicides. Oto-fure dumpsite, lies between latitude 6°27'49"N and longitude 5°36'03"E. N, the area is sloppy towards the dumpsite and crops grown include *Telfairia occidentalis*, *Capsicum annum* and *Manihot* species. The study area is characterized by tropical climate with mean annual rainfall, temperature and relative humidity of 1900 mm, 23-37°C and 89-75% respectively (NIFOR, 2018). The soils are developed from coastal plain sand parent material, with intense weathering resulting from high precipitation and temperature (Okunsebor *et al.*, 2024). Auger surface (0-15 cm) soil and *T. occidentalis* leaf samples were collected from Temboga river bank, Oto-fure dumpsite, 60 and 120 meters away from the river bank and dumpsite in 3 replicate, for laboratory analysis in the month of October 2024. Collected soil samples were air-dried, ground with wooden mortar and sieved through a 2 mm sieve, *T. occidentalis* leaf were harvested, prepared, grind in a ceramic mortar and passed via a 0.5 mm sieve. The samples (soil and *T. occidentalis* leaf), were stored in a paper bag and analyzed for physical and chemical properties in the laboratory. The soil was fractionated for Sand, silt and clay by hydrometer method (Bouyoucos, 1951), while texture was determined by textural triangle (Soil survey staff, 1999b), pH in

soil:water (1:2) suspension was measured by glass electrode pH meter (Tan, 1996), Organic carbon by wet oxidation method (Walkley and Black, 1934), available P was extracted with Bray-1 solution according to methods of Bray and Kurtz (1945), while P in the extract was developed by colorimetric method (Murphy and Riley, 1962), and measured in a spectrophotometer (Buck Scientific Model: VGP 210) at 880 nm, exchangeable bases (Ca, Mg, Na, K) were extracted with 1 N NH₄OAc, buffered at pH 7, cation exchange capacity (CEC) was calculated by the method described by Udo *et al.* (2009). Exchangeable acidity (EA) was extracted and determined by 1 M KCl and titration method respectively (Juo, 1979). Effective cation exchange capacity (ECEC) was calculated by summation of CEC and EA, while Percentage base saturation was determined by the equation given below:

$$\% \text{ Base Saturation} = \frac{CEC}{ECEC} \times 100 \dots \dots \dots (1)$$

Determination of available heavy metals

Heavy metals in Soil

Available heavy metals lead (Pb), selenium (Se), cobalt (Co), chromium (Cr), nickel (Ni), and cadmium (Cd) in soils were extracted with the diethylene triamine penta acetic acid-triethanol amine (DTPA-TEA) extractant, prepared according to the procedures described by Behera (2022) and measured using atomic absorption spectrophotometer (Buck Scientific Model: VGP 210).

Heavy metals in *Telfairia occidentalis* leaf

Heavy metals in *T. occidentalis* leaf was determined by di-acid (HNO₃ + HClO₄ at 9:4) digestion method and the digest was analyzed for available Ni, Pb, Co, Cr, Se and Cd using Atomic absorption spectrophotometer (Buck Scientific Model: VGP 210).

Statistical analysis

The relationship between the heavy metal content evaluated in soils and *Telfairia occidentalis* leaves

was determined by simple linear correlation. Data obtained from soil and *T. occidentalis* leaf analysis were statistically analyzed using the Genstat statistical package (12th edition), while Duncan multiple range tests was used to separate means at $P \leq 0.05$.

RESULTS AND DISCUSSION

Particle size distribution

Table 1 shows some physical and chemical properties of the soils studied.

Particle size distribution express the percentage sand, silt and clay fractions of a soil, soils of the experimental area have been reported to be sandy (Orhue *et al.*, 2024). Sand was the dominant soil fraction which decreased with increased distance from the river bank and dump site. Sand content in river bank soil had values of 79.13, 77.13 and 77.08% at river bank (RB), 60 meters from river bank (60 MFRB) and 120 meters from river bank (120 MFRB) respectively, while sand content had values of 78.70, 78.30 and 77.20% at dumpsite, 60 meters from dumpsite (60 MFDS) and 120 meters from dumpsite (120 MFDS), respectively. Silt content was generally low at river bank and dump site soils, silt increased slightly away from the river bank and decreased slightly away from the dumpsite, clay content at river

bank and dumpsite soils were slightly lower compare with distances away with values of 13.20 and 15.92% for river bank and dumpsite respectively, while the textural classification of the soil across both location were observed to be sandy loam. The highest value of sand content observed at the river bank could be due to sediment deposition by flowing water (Rodríguez-Rastrero *et al.*, 2023) which carries sand particles and deposits toward the river bank and dumpsite (Gupta *et al.*, 2023; Debnath *et al.*, 2023). The higher sand content in river bank and dumpsite soils as compared to soil away, found in this study is in line with the findings of Nwosu *et al.* (2018).

Some chemical properties of the soils

In the river bank soils, pH (1:2 in H_2O), of 4.73 was obtained from the river bank, while values of 5.73 and 5.40 was obtained for 60 meters away from river bank (60 MFRB) and 120 meters away from river bank (120 MFRB), while in the dumpsite soils, pH values of 5.90, 4.86 and 4.55 was obtained from dumpsite, 60 MFDS and 120 MFDS. The result indicates that the soils were acidic, while acidity decreases with increased distance away from the river bank, acidity increase with increased distance from the dumpsite, with no significance differences. Organic carbon content ranged from 1.08% at 120 MFRB to 1.85 % at river bank soil. However, or-

Table 1: Some physical and chemical properties of the soils

Location	Sand	Silt	Clay	Txt	pH	OC	Av. P	CEC	EA	ECEC	BS
	%	%	%		H_2O	%	mg/kg		cmol/kg		%
RIVER BANK SOILS											
River bank	79.13 ^a	7.67 ^a	13.20 ^a	SL	4.73 ^a	1.85 ^a	62.92 ^a	2.62 ^a	0.83 ^a	3.45 ^b	77.15 ^a
60 MFRB	77.13 ^a	8.67 ^a	14.20 ^a	SL	5.73 ^a	1.27 ^{ab}	32.83 ^b	1.75 ^b	0.99 ^a	2.70 ^b	67.80 ^b
120 MFRB	77.08 ^a	8.22 ^a	14.59 ^a	SL	5.41 ^a	1.08 ^b	13.02 ^c	1.28 ^b	3.76 ^b	5.04 ^a	27.56 ^c
SEM	3.22	1.61	1.72		0.16	0.16	36.90	0.24	0.77	0.87	8.80
SED	4.55	2.27	2.44		0.22	0.22	53.30	0.34	1.08	1.23	12.44
DUMPSITE SOILS											
Dumpsite	78.70 ^a	5.41 ^a	15.92 ^b	SL	5.90 ^a	12.10 ^{ab}	16.70 ^c	1.90 ^b	1.23 ^a	3.14 ^a	60.51 ^a
60 MFDS	78.30 ^a	4.75 ^a	16.92 ^{ab}	SL	4.86 ^a	13.23 ^a	49.00 ^b	1.08 ^a	1.00 ^a	2.08 ^a	51.92 ^b
120 MFDS	77.20 ^a	4.84 ^a	17.92 ^a	SL	4.55 ^a	11.24 ^b	54.70 ^a	1.01 ^a	0.57 ^b	1.58 ^b	63.92 ^a
SEM	1.93	0.69	1.72		0.39	1.31	11.35	0.38	0.34	0.58	8.82
SED	2.73	0.98	2.43		0.56	1.85	16.04	0.54	0.48	0.82	12.47

MFRB = meters from river bank, MFDS = meters from the dumpsite, Txt = texture, OC = organic carbon, Av. P = available phosphorus, CEC = cation exchange capacity, EA = exchangeable acidity, ECEC = effective cation exchange capacity, BS = base saturation, SL = sandy loam, SEM = standard error of means, SED = standard errors of differences of means, means with same alphabet within column are not significantly different at $p \leq 0.05$ using Duncan's multiple ranged test

ganic carbon content of 1.27 % was obtained at 60 MFRB, while in the dumpsite soils, organic carbon content ranged from 11.24 % at 120 MFDS to 13.23 % at 60 MFDS. The high organic carbon content around the river bank and dumpsite obtained in this study is in line with the study of Okebalama *et al.* (2017). The high organic carbon obtained at 60 MFRB, could be due to the deposit of organic materials from the river and use of organic manure by farmers raising vegetables and fishes around the river bank (Shao *et al.*, 2024), while higher organic carbon around the dumpsite, could have likely resulted from dumping of organic waste in the dumpsite in line with the findings of Zahra *et al.* (2024) who reported high organic carbon content in dumpsite, resulting from unsystematic solid waste dumping. Available Phosphorus (Av. P) content of 62.92, 32.83 and 13.02 mg kg⁻¹ was obtained in soils of river bank, 60 MFRB and 120 MFRB, while values of 16.70, 49.00 and 54.70 mg kg⁻¹ were obtained in soils of dumpsite, 60 MFDS and 120 MFDS, respectively. The high values of Av. P obtained in and around the river bank and dumpsite imply P sufficiency when compared to the critical level of 10-16 mg kg⁻¹ recommended by Adeoye and Agboola (1985). The high Av. P content in the river bank and soils around the river bank could be as a result of excessive use of P fertilizers by farmers around the study area in line with the findings of Mohammed *et al.* (2021), while high Av. P content around the dumpsite could be due to P mineralization from disposed waste, in line with the report of Okafor and Obaze (2024), who concluded that waste disposal sites have positive effects on soil fertility. High Av. P could also be due to the higher percent organic carbon content obtained in both river bank and dumpsite location, as organic matter have been reported to supply 75% of organic Phosphorus, in line with the studies of Orhue *et al.* (2024). The cation exchange capacity (CEC) and percentage base saturation (% BS) was observed to be the highest at the river bank while exchangeable acidity (EA) and effective cation exchange capacity (ECEC) were highest at 120 MFRB. The CEC and % BS had highest values of 2.62 cmolkg⁻¹ and 77.15% at the river bank soil respectively, which implies higher capacity of the river bank soils to hold and exchange basic cations. However, the CEC val-

ues of 1.72 Cmol kg⁻¹ and 1.28 Cmol kg⁻¹ obtained for 60 MFRB and 120 MFRB soils respectively were in line with CEC results reported by Orhue *et al.* (2024). The ECEC values obtained from the river bank, dumpsite and their surrounding soils could be due to high levels of exchangeable acidity obtained in the soils as reflected in the results of the study. However, the values of ECEC obtained in the study were below 15 cmol kg⁻¹ critical values reported by Udo *et al.* (2009) for tropical soils. The CEC, ECEC and base saturation status of a soil gives an insight on the fertility status (Onwuka and Ani, 2023). Orhue *et al.* (2024) have also reported that ECEC, could be used to evaluate the fertility status of a given soil.

Heavy metals content in soils around river bank and dump site

The heavy metal content in soils of river bank and dump site and are shown in Table 2.

The result showed that heavy metal (Ni, Pb, Co, Cr, Se and Cd) levels varied with distances away from river bank and dumpsite. In river bank soils, Ni levels were below the critical level of 10-1000 mgkg⁻¹ reported by Allaway (1968). However, values of 1.36, 0.22 and 1.37 mg Ni kg⁻¹ was obtained from river bank, 60 meters from river bank (60 MFRB) and 120 meters from river bank (120 MFRB), while values of 3.13, 1.28 and 1.61 mg Ni kg⁻¹ were obtained at dumpsite, 60 meters from dumpsite (60 MFDS) and 120 meters from dumpsite (120 MFDS) respectively. This result indicated Ni contamination but below toxicity level around the river bank and dumpsite. The low Ni content obtained could be due to Ni absorption by crops grown in the area, contrary to the findings of Fayad *et al.* (2013) who reported higher level of Ni in river bank soils. The higher level of Ni reported by Fayad and his colleagues could be due to deterioration of the river bank over the years due to poor farming practices and industrial pollution. The Pb and Se content were higher at the river bank and dumpsite soils compared to the surrounding soils, the Pb and Se had values of 0.03, 0.02, 0.01 mg Pb kg⁻¹ and 0.06, 0.02, 0.01 mg Se kg⁻¹ for river bank, 60 MFRB and 120 MFRB, while values of 0.03, 0.02 and 0.02 mg Pb kg⁻¹ and 0.07, 0.05

and 0.03 mg Se kg⁻¹ were obtained for dumpsite, 60 MFDS and 120 MFDS respectively. Although, these values of Pb and Se were below critical values (2-200 mg Pb kg⁻¹ and 0.01-2 mg Se kg⁻¹) reported by Allaway (1968) and Dhillion and Dhillion (2009). The higher values obtained at the river bank and dumpsite soils implies that the Pb and Se contamination could be due to enrichment of the surrounding soils by the river water and waste pile respectively. Se levels in the river bank, dumpsite and surrounding soils is tending towards the 2 mg Se kg⁻¹ upper limit and could reach toxic level in a short time. The higher Pb and Se content around the river bank found in this study align with the findings of Fayad *et al.* (2013) who reported higher heavy metal

content around a river bank. Co and Cd content varied with distances from river bank and dumpsite, but were below critical values (1-40 mg kg⁻¹ for Co and 0.01-0.7 mg kg⁻¹ for Cd) reported by Allaway (1968). These values of Co and Cd content obtained could be due to anthropogenic activities around the river bank and dumpsite. This result of decreased Co content with increase distance away from the river bank found in this study is in line with the finding of Ayeni *et al.* (2010). The Cr content decreased with increased distance from the river bank and dumpsite, but the soils could be said to be low in Cr when compared with the critical range of 5-3000 mg Cr kg⁻¹ (Allaway, 1968). The result of this study align with the findings of Nnamonu *et al.* (2015) also reported similar

Table 2: Heavy metals content in River bank and Dumpsite soils in mg/kg

Location	Ni	Pb	Co	Cr	Se	Cd
RIVER BANK SOIL						
River bank	1.36 ^a	0.03 ^a	0.16 ^a	0.46 ^a	0.06 ^a	0.01 ^a
60 MFRB	0.22 ^b	0.02 ^a	0.16 ^a	0.21 ^b	0.02 ^b	0.01 ^a
120 MFRB	1.37 ^a	0.01 ^a	0.05 ^b	0.22 ^b	0.01 ^b	0.01 ^a
SEM	0.41	0.01	0.07	0.14	0.00	0.01
SED	0.58	0.01	0.09	0.19	0.01	0.01
DUMPSITE SOILS						
Dumpsite	3.13 ^a	0.03 ^a	0.03 ^b	1.02 ^a	0.07 ^a	0.01 ^a
60 MFDS	1.28 ^b	0.02 ^a	0.11 ^a	0.36 ^b	0.05 ^b	^B DL
120 MFDS	1.61 ^b	0.02 ^a	0.06 ^b	0.08 ^b	0.03 ^c	0.00 ^a
SEM	1.26	0.01	0.04	0.27	0.02	0.01
SED	1.78	0.01	0.06	0.38	0.02	0.01

MFRB = meters from river bank, MFDS = meters from dumpsite, BDL= below detectable limit, SEM = standard error of means, SED = standard, errors of differences of means, means with same alphabet within column are not significantly different at $p \leq 0.05$ using Duncan's multiple range test

Table 3: Heavy metals content in *Telfairia occidentalis* leaf in mg/kg

Location	Ni	Pb	Co	Cr	Se	Cd
RIVER BANK SOILS						
River bank	2.56 ^a	0.19 ^b	0.46 ^b	0.13 ^c	0.31 ^a	BDL
60 MFRB	0.67 ^b	0.39 ^a	0.69 ^b	0.76 ^a	0.16 ^b	0.03 ^a
120 MFRB	1.29 ^c	0.03 ^c	1.51 ^a	0.22 ^b	0.24 ^a	0.02 ^a
SEM	0.81	0.14	0.66	0.37	0.05	0.012
SED	1.14	0.19	0.93	0.52	0.08	0.02
DUMPSITE SOILS						
Dumpsite	31.50 ^a	0.27 ^c	1.21 ^a	8.92 ^a	1.46 ^a	BDL
60 MFDS	13.00 ^b	0.46 ^a	1.59 ^a	4.36 ^{ab}	1.42 ^b	BDL
120 MFDS	12.50 ^b	0.40 ^a	1.16 ^a	1.77 ^a	0.67 ^b	BDL
SEM	13.02	0.16	0.29	1.45	0.14	-
SED	18.41	0.23	0.41	2.04	0.20	-

MFRB = meters away from the dumpsite, MFDS = meters from dump site, BDL= below detectable limit, SEM = standard error of means, SED = standard errors of differences of differences of means, means with same alphabet within columns are not significantly different at $p \leq 0.05$ using Duncan's multiple range test

Table 4: Relationship between heavy metal in soil and *T. occidentalis* leaf

Location	Ni	Pb	Co	Cr	Se	Cd
RIVER BANK SOILS						
River bank	0.999*	0.878	0.999*	0.955*	-0.711	-
50 MFRB	0.999*	-0.947	0.668	0.176	0.204	-0.982
100 MFRB	0.049	0.974	-0.634*	0.999*	-0.577	1.000***
DUMPSITE SOILS						
Dumpsite	0.99*	0.04	0.76	0.99*	0.76	0.00
60 MFDS	0.98	0.87	0.62	0.90	-0.61	0.00
120 MFDS	0.99	-0.27	0.72	-0.12	-0.53	0.00

MFRB = meters from river bank, MFDS = meters from the dumpsite, * = significant at $p \leq 0.05$, *** = significant at $p \leq 0.001$

trend of Cr concentration in soils around river bank than soil away from river bank.

Heavy metals in fluted pumpkin (*Telfairia occidentalis*) leaf cultivated around river bank and dumpsites

The result of heavy metal content in fluted pumpkin (*Telfairia occidentalis*) leaf is presented in Table 3. The result shown that Ni, and Se content of *T. occidentalis* leaf grown around the river bank and dumpsite, decreased with increased distance from the river bank and dumpsite, while Cd was below 0.2–0.8 mg kg⁻¹ (detectable limit) in *T. occidentalis* leaf grown around at river bank and all dumpsite locations. Low Cd content (0.05 mg Cd kg⁻¹) in vegetables grown around river bank soil with some below detectable limit (BDL) has been reported (Kihampa and Mwegoha, 2010). Pb, Co and Cr content in *T. occidentalis* leaf were inconsistent with distances away from river bank and dumpsite; however Cr decreased with increased distance from the dumpsite. *T. occidentalis* leaf grown at river bank, 120 MFRB and all locations around the dumpsite contains excessive content of Ni as against the 1.0 mg Ni kg⁻¹ permissible limit reported by Allaway (1968), while Ni buildup was observed in *T. occidentalis* grown at 60 MFRB, these excessive levels of Ni in the *T. occidentalis* leaf may pose health risk to consumers who regularly consume the *T. occidentalis* leaf grown around the river bank and dumpsite. *T. occidentalis* leaf grown around the river bank and dumpsite, were found to contain toxic levels of Co above permissible level of 0.05–0.5 mg Co kg⁻¹ reported by Allaway (1968) which could result in Co toxicity, and can pose

health risk to food chain and humans who regularly consume *T. occidentalis* grown around the river bank and dumpsite. The result also showed that Pb, Cr and Se were building up in *T. occidentalis* leaf grown around the river bank, while *T. occidentalis* leaf grown around dumpsite contains toxic levels of Cr, compared to permissible levels reported in literatures, the result corroborate the findings of Kihampa and Mwegoha (2010) who reported high Pb, Cr and Se content in vegetables grown around river banks. This result of toxic and buildup levels of heavy metals in the *T. occidentalis* leaf, found in this study could be due to bioavailability of from surrounding soils (Osemudiamen *et al.*, 2023; Oladeji and Saeed, 2015).

Relationship between heavy metals in soils and *Telfairia occidentalis* leaf

Table 4 shows the relationship between heavy metals in soils and *T. occidentalis* leaf.

The result showed that there were positive and negative relations between heavy metal content in soils and *T. occidentalis* leaf in the river bank and dumpsite soils. Significant correlation was observed for Ni at river bank (0.999 $p \leq 0.05$) and 60 MFRB (0.999 $p \leq 0.05$). Significant correlation was also observed for Co and Cr content at river bank (0.999 $p \leq 0.05$ and 0.955 $p \leq 0.05$) and 120 MFRB (-0.634 $p \leq 0.05$ and 0.999 $p \leq 0.05$) respectively, while Cd was highly significance at 120 MFRB (1.000 $p \leq 0.001$), however, significant positive relationship between soil and *T. occidentalis* leaf content was observed for Ni (0.990 $p \leq 0.05$) and Cr (0.990 $p \leq 0.05$) at dumpsite location. This implies that increase in heavy metal

levels in soil will lead to significant increase in the amount of bioavailable heavy metals in *T. occidentalis* leaf except for Co at 120 MFRB where antagonistic relationship was obtained. Emurotu and Onianwa (2017) also reported significant relationship between heavy metals content in soils and *T. occidentalis*.

CONCLUSION

Soils along the river bank, dumpsite and adjacent areas contains potentially toxic levels of Se (0.06 mg kg⁻¹ for riverbank, 0.07 mg kg⁻¹ for dumpsite) and Cd (0.01 mg kg⁻¹ for riverbank and dumpsite), although all the heavy metal studied were found to be highest at the river bank and dumpsite soil except Cd, suggesting that soil enrichment may be due to river overflow and waste pile in dumpsite, and could potentially reach toxic levels over time if not properly managed. Bioaccumulation of Ni (2.56 mg kg⁻¹ for riverbank, 31.50 mg kg⁻¹ for dumpsite) and Cr (0.13 mg kg⁻¹ for riverbank, 8.92 mg kg⁻¹ for dumpsite) in *T. occidentalis* leaf, has reached toxic levels, while Pb, Cd, Se and Co accumulation are at potential toxic level (building up) in *T. occidentalis* leaf grown around river bank and dumpsite with values approaching or exceeding the critical levels, which could be due to bioavailability of these heavy metals in surrounding soils. The findings suggests that the consumption of *T. occidentalis* leaves grown around Temboga river bank and Oto-fure dumpsites may pose risk to the food chain and human health, emphasizing the need for further research and mitigation strategies to reduce heavy metal enrichment in the soils.

The significant correlation observed between Ni (0.999 $p \leq 0.05$ for riverbank, 0.990 $p \leq 0.05$ for dumpsite), and Cr (0.995 $p \leq 0.05$ for riverbank, 0.990 $p \leq 0.05$ for dumpsite) concentration in soil and *T. occidentalis* leaf indicate that increased levels of these heavy metals in soil can lead to elevated accumulation in the crop. It is also recommended that cultivation and consumption of *T. occidentalis* leaf grown at the Oto-fure dumpsite and Temboga river bank, be discouraged as it has been found to contain toxic levels of Ni and Cr, with elevated lev-

els Pb, Cd, Se and Co.

ACKNOWLEDGEMENTS

We wish to appreciate Prof. E. R. Orhue of Department of Soil Science and Land management, University of Benin, Nigeria, for designing and supervising this project, we are also grateful to all the vegetable farmers in Temboga river bank and Oto-fure dumpsite, Benin City for their approval to collect soil and *T. Occidentalis* leaf samples from their field, we also wish to acknowledge and congratulate students of 2023/2024 soil chemistry project sect, in the Department of Soil Science and Land management, for their diligence during the laboratory analysis of the soil and *T. occidentalis* leaf samples.

REFERENCES

- Acharya, S. (2024). Heavy Metal Contamination in Food: Sources, Impact, and Remedy. In Food Safety and Quality in the Global South. Springer Nature Singapore, Pp: 233-261.
- Adefarati, O., Ahmed O. O., Zainab A. S., Rasheedat O. A. and Sunday, A. K. (2024). Heavy metal contamination in soils, water, and food in Nigeria from 2000–2019: A systematic review on methods, pollution level and policy implications. *Water, Air and Soil Pollution*, 235 (9): 586.
- Adeoye, G. O. and Agboola, A. A. (1985). Critical levels for soil pH, available P, K, Zn and Mn and maize ear-leaf content of P, Cu and Mn in sedimentary soils of South-Western Nigeria. *Fertilizer Research*, 6: 65–71.
- Akande, M. G., Sanni, F. S. and Enefe, N. G. (2021). Assessment of the concentrations and health risk of some heavy metals in cowpea (*Vigna unguiculata*) in Gwagwalada, Nigeria. *Drug and Chemical Toxicology*, 44 (5): 518-523.
- Allaway, W. H. (1968). Agronomic controls over environmental cycling of trace elements. *Advances in Agronomy*, 20: 235–274.
- Amadi, C. N., Frazzoli, C. and Orisakwe, O. E. (2020). Sentinel species for biomonitoring and biosurveillance of environmental heavy metals in Nigeria. *Journal of Environmen-*

- tal Science and Health*, 38 (1): 21-60.
- Ayeni, O. O., Ndakidemi, P. A., Snyman, R. G. and Odendaal, J. P. (2010). Metal contamination of soils collected from four different sites along the lower Diep River, Cape Town, South Africa. *International Journal of Physical Sciences*, 5 (12): 2045-2051.
- Ayoola, A. J. (2021). Heavy metals in soils: Trace metals and metalloids in soils and their bioavailability. *Environmental Pollution*, Pp 50–102.
- Behera, R. D. (2022). Soil and Plant Analysis. NIPA GENX Electronic Resources and Solutions P. LTD. New Delhi-110 034, Pp: 81-83.
- Bhagwat, V. R. (2019). Safety of water used in food production. In R. L. Singh and S. Mondal (Eds.), Food Safety and Human Health, Elsevier Inc. (first), Pp: 219–247.
- Bouyoucos, G. J. (1951). Hydrometer method improved for making particle size analysis of soils. *Agronomy Journal*, 54: 464-465.
- Bray, R. H. and Kurtz, L. T. (1945). Determination of Total Organic and Available Forms of Phosphorus in Soils. *Soil Science*, 59: 39-45.
- Debnath, J., Sahariah, D., Saikia, A., Meraj, G., Nath, N., Lahon, D., Annayat, W., Kumar, P., Chand, K., Singh, S. K. and Kanga, S. (2023). Shifting sands: assessing bankline shift using an automated approach in the Jia Bharali River, India. *Land*, 12 (3): 703.
- Dhillion, S. K. and Dhillion, K. S. (2009). Phytoremediation of selenium-contaminated soils: the efficiency of different cropping systems. *Soil Use and Management*, 25 (4): 441-453.
- Edelstein, M. and Ben-Hur, M. (2018). Heavy metals and metalloids: Sources, risks and strategies to reduce their accumulation in horticultural crops. *Scientia Horticulturae*, 234: 431-444.
- Emurotu, J. E. and Onianwa, P. C. (2017). Bioaccumulation of heavy metals in soils and selected food crops cultivated in Kogi State, North central, Nigeria. *Environmental System Research*, 6 (21): 22-31.
- Fayad, N., Al-Noor, T. H. and Al-Noor, N. H. (2013). Analysis and assessment of essential toxic heavy metals, pH and EC in Ishaqi River and adjacent soil. *Advances in Physics Theories and Applications*, 16: 25-37.
- Friedlova, M. (2010). The influence of heavy metals on soil biological and chemical properties. *Soil and Water Research*, 5 (1): 21–27.
- Gupta, L. K., Pandey, M., Raj, P. A., Shukla, A. K. (2023). Fine sediment intrusion and its consequences for river ecosystems: A review. *Journal of Hazardous, Toxic, and Radioactive Waste*, 27 (1): 04022036.
- Islam, M. S., Ahmed, M. K., Al-mamun, M. H. Islam, S. A. M. (2019). Sources and Ecological Risks of Heavy Metals in Soils Under Different Land Uses in Bangladesh. *Pedosphere*, 29 (5): 665–675.
- Juo, A. S. R. (1979). Selected Methods for Soil and Plant Analysis. Manual Series No.1, IITA, Ibadan.
- Kihampa, C. and Mwegoha, W. J. S. (2010). Heavy metals accumulation in vegetables grown along the Msimbazi river in Der es Salaam, Tanzania. *International Journal of Biological and Chemical Sciences*, 4 (6): 63-71.
- Lissy, A. and Madhu, D. (2020). Transport and Remediation of Subsurface Contaminants, American Chemical Society, Washington DC, USA, Pp 108-120
- Mohammed, S. B., Dzidzienyo, D. K., Yahaya, A., Umar, L. M., Ishiyaku, M. F., Tongoona, P. B. and Gracen, V. (2021). High Soil Phosphorus Application Significantly Increased Grain Yield, Phosphorus Content but not Zinc Content of Cowpea Grains. *Agronomy*, 11 (4):802.
- Murphy, J. and Riley, J. P. (1962). A modified single solution method for the determination of phosphate in natural waters. *Analytical Chemistry ACTA*, 27: 31-36.
- NIFOR (Nigerian Institute for Oil Palm Research) (2018). Weather data (Temperature, Rainfall, Relative Humidity): 1993-2018. Nigerian Institute for Oil Palm Research Main Station, Benin-City, Nigeria.
- Unabueze, D. N., Helen, O., Taghi, M. and Chinenye, U. (2023). Bioaccumulation for heavy metal

- removal: a review. *SN Applied Sciences*, 5 (5): 125.
- Nnamonu, L. A., Ogidi, O. A. and Eneji, I. S. (2015). Analysis of heavy metals content of fluted pumpkin leaves cultivated on the south bank of river benue, Nigeria. *Food Science Quality Manage*, 39: 1-8.
- Nwosu, O. C., Osadeke, V. E. and Ohaeri, J. E. (2018). Forms and profiles distribution of phosphorus soils along a toposequence of Amaoba-Ime, Ikwuano LGA, Abia State, Nigeria. *Nigerian Agricultural Journal*, 49 (1): 101-108.
- Okafor, O. C. and Obaze, W. O. (2024). Environmental impact of waste disposal sites on physicochemical properties of soil in Enugu State, Nigeria. *Arabian Journal of Geosciences*, 17 (6): 190.
- Okebalama, C. B., Igwe, C. A. and Okolo, C. C. (2017). Soil organic carbon levels in soils of contrasting land uses in Southeastern, Nigeria. *Tropical and Subtropical Agroecosystems*, 20 (3): 493-504.
- Okunsebor, F. E., Ogbemudia, I. and Okolie, S. I. (2024). Characterization and fertility capability classification of some soils in the rain forest zone of Edo state, Nigeria. *Pantnagar Journal of Research*, 22 (1): 14-25
- Oladeji, S. O. and Saeed, M. D. (2015). Assessment of cobalt levels in wastewater, soil and vegetable samples grown along Kubanni stream channels in Zaria, Kaduna State, Nigeria. *African Journal of Environmental Science and Technology*, 9 (10): 765-772.
- Onwuka, O. and Ani, J. O. (2023). Evaluation of Fertility Status of Soils of Different Parent Materials used for Arable Crop Production in Enugu State, Nigeria. *International Journal of Agriculture and Rural Development*, 26 (2): 6754-6758.
- Orhue, E. R., Bakare, A. O. and Emomu, A. (2024). Zinc, Cobalt and Chromium status in soils under citrus and thick plantation, University of Benin, Edo State, Nigeria. *Journal of Sustainable Development*, 22 (1): 88-98.
- Osamudiamen, E., Grace, O. and Ozekeke, O. (2023). Nickel and Vanadium in edible fruits and vegetables, potential health risk or not? *Journal of Science and Technology Research*, 5 (2): 92-101.
- Pandit, D., Haque, M. M., Bhuyan, M. S., Harun-Al-Rashid, A., Barman, P. P., Roy, R., Bishwajit, Sarker, B., Saifulah, M. K., Kunda, M. (2024). A comprehensive scenario of heavy metals pollution in the rivers of Bangladesh during the last two decades. *Environmental Science and Pollution Research*, 32: 10240-10257.
- Rahi, A. A., Younis, U., Ahmed, N., Ali, M. A., Fahad, S., Sultan, H., Zarei, T., Danish, S., Taban, S., El Enshasy, H. A. and Tamunaidu, P. (2022). Toxicity of Cadmium and nickel in the context of applied activated carbon biochar for improvement in soil fertility. *Saudi Journal of Biological Sciences*, 29 (2): 743-750.
- Rodríguez-Rastrero, M., Suárez, C. E., Ortega, A., Cuevas, J. and Fernández, R. (2023). Geochemical Anomalies in Soils and Surface Waters in an Area Adjacent to a Long-Used Controlled Municipal Landfill. *Sustainability*, 15 (23):16280.
- Shao, G., Xu, Y., Zhou, J., Tian, P., Ai, J., Yang, Y., Zamanian, K., Zeng, Z. and Zang, H. (2024). Enhanced soil organic carbon stability in rhizosphere through manure application. *Soil and Tillage Research*, 244: 106223.
- Skuza, L., Szuæko-Kociuba, I., Filip, E. and Bożek, I. (2022). Natural molecular mechanisms of plant hyperaccumulation and hypertolerance towards heavy metals. *International Journal of Molecular Sciences*, 23 (16): 9335.
- Soil survey staff (1999b). Soil Taxonomy: A basic system of soil classification for making and interpreting soil survey. 2nd edition. Natural Resource conservation Service. US Department of Agriculture Handbook, 436 p.
- Tan, K. M. (1996). Soil sampling preparation and analysis. In: African Research Review (Volume 3, 23-33). Mercel Dekkes Inc., New York, Basel, Hong Kong
- Toheeb, L. J., Isa, O. E., Prince, E. N., Taiwo, E. F., Opeyemi, Y. A., Ibrahim, O. O., Faruq, E. A., Adejuwon, A. M., Zechariah, O. O. and

- Ogunkoya, O. I. (2024). *Environmental Monitoring and Assessment*, 196 (2): 109.
- Udo, E. J., Ogunwale, J. A., Fagbami, A., Ano, A. O. and Esu, I. E. (2009). Manual of soil, plant and water resources. Sibon books limited, 177 p.
- Wang, Y., Zhang, Z., Li, Y., Liang, C., Huang, H. and Wang, S. (2024). Available heavy metals concentrations in agricultural soils: Relationship with soil properties and total heavy metals concentrations in different industries. *Journal of Hazardous Materials*, 471: 134410.
- Xia, X., Wu, S., Zhou, Z. and Wang, G. (2021). Microbial Cd (II) and Cr (VI) resistance mechanisms and application in bioremediation. *Journal of Hazardous Materials*, 401: 123685.
- Zahra, B., Begum, F., Seung-Jun, W. O. O. and Min-Jae, J. U. N. G. (2024). Impacts of Unsystematic Solid Waste Dumping on Soil Properties and Climate Change. *Journal of Wellbeing Management and Applied Psychology*, 7 (3): 31-42.

Received: April 23, 2025

Accepted: July 08, 2025