

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 24(1) January-April 2026]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology
Pantnagar, U.S. Nagar; Uttarakhand, Website : gbpuat.res.in/PJR



ADVISORY BOARD

Patron

Prof. Shivendra Kumar Kashyap, Ph.D., Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. S. K. Verma, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. D. Kumar, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Lokesh Varshney, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Subhash Chandra, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Ramesh Chandra Srivastava, Ph.D., Dean, College of Basic Sciences & Humanities, G.B.P.U.A.T., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W.P. Carey Business School, Arizona State University, U.S.A.

Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Indu S. Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agri. & Tech., Pantnagar, India

Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

Nikitin Georgy, Ph.D., Vice-Rector cum Professor St. Petersburg State University of Veterinary Medicine St. Petersburg, Russia

Sanjay Jambhulkar, Ph.D., Ex-Scientific Officer 'H' Head Mutation Breeding Section Bhabha Atomic Research Centre, Mumbai Professor, Homi Bhabha National Institute, Mumbai

Vijay Kubsad, Ph.D., Adjunct Professor Washington State University USA

Kiran Tota-Maharaj, Ph.D., Professor Royal Agricultural University (RAU) Cirencester, Gloucestershire GL76JS England, UK

Ravindra P. Veeranna, Ph.D., Dean, Xavier University, College of Veterinary Sciences, Aruba

Javed Akhtar, Ph.D., University of Technology and Applied Sciences, Muscat

Basant Maheshwari, Ph.D., Western Sydney University, Australia

Vijay P. Singh, Ph.D. Professor Emeritus, Texas A&M University College station, Texas, USA

Sanjay Sharma, Ph.D. Director, Uttarakhand Council of Biotechnology

Krishna Nemali, Ph.D. Controlled Environment Agriculture Extension Specialist

Editor-in-Chief

K.P. Raverkar, Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Managing Editor

S.K. Guru, Ph.D., Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head

Brijesh Dumka, Developer & Programmer

CONTENTS

Indian mustard: Phytoremediation tool for soil health management (Review)	1
M.L. DOTANIYA, C.K. DOTANIYA, KULDEEP KUMAR, JAKIR HUSSAIN, R.K. DOUTANIYA, H.M. MEENA and MANJU LATA	
From profile to problems: A study on socio-economic status and constraints of rural women in dairy farming	14
ADITI PATHAK and ARPITA SHARMA KANDPAL	
Diversification and intensification in rice–wheat cropping system for enhancing productivity under Tarai zone of Uttarakhand	20
PESALA SANTHA KUMAR, DINESH KUMAR SINGH, SUBHASH CHANDRA, ROHITASH SINGH and AJEET PRATAP SINGH	
Sensor based nitrogen application impact on productivity, economics and nutrient uptake in maize	25
AMIT BHATNAGAR and AJAY KUMAR	
Association of germination and seedling vigour attributes with green forage yield in oat (<i>Avena sativa</i> L.) genotypes through correlation and path coefficient analysis	31
DEVYANI RAMOLA, AKRITI BALLABH SANSKRITI JOSHI, MANISH MEHRA and BIRENDRA PRASAD	
Genetic divergence and yield trait impact in French bean (<i>Phaseolus vulgaris</i> L.)	43
BHAWANA MAMGAIN, B.P. NAUTIYAL, S.C. PANT, VAISHALI THAPA, HEIGRUJAM RIYASANA and MAYANK	
Arabic-gum based edible coatings preserving the post-harvest quality of Guava fruits	48
ESHA TEWARI and OMVEER SINGH	
Efficacy of catechol and plant growth regulators on the vegetative growth and flowering of African Marigold (<i>Tagetes erecta</i> L. cv. Pusa Narangi Gainda)	59
SURYANSH SRIVASTAVA, AFSAR AHMAD, MD. RAMJAN and KHAN BILAL MUKHTAR AHMED	
Host response of diverse pulse genotypes to root-knot nematode (<i>Meloidogyne incognita</i>)	67
SAROJ YADAV, EKTA RATHORE and ANIL KUMAR	
Effect of egg size on egg quality traits, fertility, hatchability and day-old chick weight	75
ISLAM UDDIN SHEIKH, AZMAT ALAM KHAN, ARNAB JYOTI KALITA, SHEIKH ADIL HAMID, IRFAN AKRAM BABA, HENNA HAMADANI, BUSHRA ZAFFER and INSHA AFZAL	
Pathomorphological studies on verminous pneumonia in a Gaddi goat	81
SWASTI SHARMA, SONALI MISHRA and R.D. PATIL	
Pathological findings in congestive heart failure in dog: A case report	85
MUNISH BATRA and STUTI SINGH	

Histopathological and haematological evaluation of the ameliorative effect of piperine on quinalphos-induced subacute toxicity in Swiss Albino Mice PREETI SINGH, VINOD KUMAR, PREETI BAGRI and DEEPIKA LATHER	89
Phytochemical screening and antioxidant activities of locally available medicinal plants PREETI BAGRI, VIJEYTA TIWARI, ARCHANA LOHIYA and VINOD KUMAR	99
Natural Farming Inputs (NFIs) as sustainable alternatives for enhancing the growth and development of Chickpea (<i>Cicer arietinum</i> L.) SHAIFALI and A.K. SHARMA	108
Integrating Bayesian Calibration, Hierarchical Spatial Regression, and Structural–Temporal Models to Explain and Predict Wheat Yield in the Tarai–Bhabar Belt of Uttarakhand, India NEHA BHATT and VINOD KUMAR	114
Phytochemical characterization and immuno-antioxidant potential of <i>Moringa oleifera</i> leaf extract in chicken lymphocytes ANAMITRA RUJ, SONU AMBWANI and LATA PALIWAL	128
Biochemical changes associated with triclosan exposure in the freshwater fish <i>Cyprinus carpio</i> MADHU SHARMA, YASHPAL, RIJUL RANA, DEEKSHA, DEEPIKA THAKUR and TARANG KUMAR SHAH	140
Assessment of awareness among respondents towards Pradhan Mantri Jan Dhan Yojana: A study in Haryana MEGHA GOYAL, SUMAN GHALAWAT, MINAKSHI, ATUL DHINGRA and MINI	148
Youth perception and constraints in agri-entrepreneurship: A factor analytic study JYOTI, SUMAN GHALAWAT, MEGHA GOYAL, NEELAM KAUSHAL, SUNIL KUMAR and VIKRANT HOODA	156
Design of Object Detection Model Using Deep Learning for Autonomous Driving in Adverse Weather Conditions: A GAN-Based Restoration and Depth-Aware Fusion Framework MUMUKSHU BHATT, SUBODH PRASAD, RAJESH SINGH, ASHOK KUMAR and POOJA TAMTA	165
A novel mechanical purification approach for <i>Grewia optiva</i> Fibre-Reinforced epoxy composites: Effect of fibre geometry on structural performance DEEPA SINGH, NEERAJ BISHT, TARUN KUMAR SAXENA and KAMAL RAWAT	172
Evaluating Long-Term rainfall and temperature trends across the Shipra River basin POONAM SINGH, MOHAN LAL, DHEERAJ KUMAR, P. K. SINGH and GOVIND VERMA	184
Rainfall-runoff modelling using soft computing techniques for various watersheds of Madhya Pradesh, India SEEMA KUSHWAHA and PRAVENDRA KUMAR	196
Prevalence and associated risk factors of functional constipation among the selected adult population of Pantnagar, U.S. Nagar, Uttarakhand, India JYOTI SINGH, NEETU DOBHAL, ARCHANA KUSHWAHA, N.C SHAHI and SABBU SANGEETA	204
Tracking nutritional transitions: A comparative study of child malnutrition (0–5 years) trends across districts of Madhya Pradesh SHALINI CHAKRABORTY	212
Determinants influencing tourist selection of rural tourism destinations using Garrett Ranking Technique: Evidence from Supi village, Uttarakhand NEHA PANDEY and KAMAL UPRETI	221

Effect of egg size on egg quality traits, fertility, hatchability and day-old chick weight

ISLAM UDDIN SHEIKH*, AZMAT ALAM KHAN, ARNAB JYOTI KALITA, SHEIKH ADIL HAMID, IRFAN AKRAM BABA, HENNA HAMADANI, BUSHRA ZAFFER and INSHA AFZAL

Division of Livestock Production and Management, Faculty of Veterinary Sciences and Animal Husbandry, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, Srinagar (UT of Jammu and Kashmir)

**Corresponding author's email id: sheikhiu@gmail.com*

ABSTRACT: An experiment was conducted on 500 Vanaraja eggs of different sizes viz., T₁ (40-45g), T₂ (45-50g), T₃ (50-55g), T₄ (55-60g) and T₅ (60-65g) to study the egg quality traits, fertility, hatchability and weight of day-old chicks in an organized farm. A minimum of 100 eggs per group were selected and incubated under optimum temperature, humidity and other conditions required for incubation was maintained in setter. Ten eggs from each group were selected to study different external and internal quality parameters. Candling was performed on 18th day of incubation to determine fertility of the eggs and fertile eggs were transferred to hatcher. Fertility (%) ranged from 71.30±4.67 per cent in T₁ to 84.00±7.00 per cent in T₄ group with significant (P≤0.05) differences and hatchability was also significant (P≤0.05) among the groups. The body weight of day-old chicks ranged from 27.67±1.49 in T₁ to 42.34±0.85 g in T₅ groups with significant (P≤0.05) differences among the groups. The dead germ (%) was significantly (P≤0.05) higher in T₁ (7.33±0.67) group than all other groups. The dead -in- shell (%) was significantly (P≤0.05) lowest in T₃ (14.50±1.50) and highest in T₁ group (19.25±1.15). The egg specific gravity (mg/dL) was ranged from 1.073±0.02 to 1.092±0.03. The albumen weight ranged from 24.31±0.87 to 38.59±1.12g with significant (P≤0.05) differences. The Haugh unit scores among the groups differed significantly (P≤0.05). From the present study it is concluded that the size of egg has significant effect on fertility, hatchability and dead-in-shell of Vanaraja parent stock eggs. The body weight of day-old chicks significantly (P≤0.05) increased linearly as the egg weight increased.

Key words: Chick weight egg quality, Fertility, Hatchability, Vanaraja

The size of eggs plays an important role in hatchability of eggs. The body weight of day-old chick is also an important factor for growth rate of the birds reared in the farms. The higher body weight at hatch grows faster than their counterpart of lower body weight. The performance of the chicken in terms of hatchability and chick hatch weight has been found closely associated to egg weight (Mbajiorgu and Ramaphala, 2026). If the birth weight i.e., weight of day old chick is optimum, then the chicks will grow at faster rate in the subsequent rearing period besides genetic, nutrition and other managerial factors. Fertility and hatchability plays significant role in profitability of the hatchery enterprise for small and medium-sized stakeholders (Kuda, 2023; Rathnayake *et al.*, 2024). The flock age plays a pivotal role in both egg production and the hatchability of broiler eggs which is a multifaceted trait that changes with time (Peric *et al.*, 2022). The supply of day-old chick is highly influenced by

fertility and hatchability of egg. Fertility refers to the percentage of incubated eggs that are fertile while hatchability is the percentage of fertile eggs that hatch. Therefore, it is important to understand the factors responsible or influence the fertility and hatchability of eggs. Egg size had a significant effect on fertility and hatchability traits. The medium egg size group had the highest percentages of hatchability and fertility, followed by the small and large sized egg (Kumar *et al.*, 2025). There are many factors contributing to the failure of a fertile egg to hatch, that includes lethal genes, insufficient nutrients in the egg and exposure to the conditions that does not meet the needs of the developing embryo (Mabotha *et al.*, 2025). The egg weight, shell thickness, porosity, shape index and the consistency of the contents are most influential egg parameters that influence hatchability. Although breed has little effect on fertility and hatchability of poultry eggs but higher fertility and hatchability in light breeds

of chicken have been reported (Gandhimathi *et al.*, 2020). In most of the cases embryonic mortality that occurs during the incubation period, is related to inadequate incubation conditions rather than poor egg quality and an imbalanced diet of the breeder flock (Agbehadzi *et al.*, 2024). Egg weight directly affects hatchability, embryonic mortality, hatching weights, and subsequent performance of chicks (Fathi *et al.*, 2022; Mbajiorgu and Ramaphala, 2026). Therefore, the present study was carried out to find out the ideal hatching egg size for better egg quality traits, fertility, hatchability, moisture loss and chick weight of Vanaraja chicken.

MATERIALS AND METHODS

A total of five hundred (500) hatching eggs comprising of one hundred (100) eggs of each size category: T₁ (40-45g), T₂ (45-50g), T₃ (50-55g), T₄ (55-60g) and T₅ (60-65g) were selected which were clean and defect free, oval in shape and set into five trays. All eggs were weighed by Denver XP-300 electronic balance corrected to 0.1 grams. Average egg weight of each tray having 100 hatching eggs (experimental eggs) was calculated. These five trays of each egg size category were randomly placed with broad end up to the top, middle and lower part of the setter machine (model ISI, having capacity of 2000 eggs) to account for possible environmental difference caused by position in incubator. The setter was operated at a temperature of 99.5°F and a relative humidity of 55-60% during the first 18 days of incubation and eggs were turned after every three-hour interval. Candling of eggs performed on 18th day of incubation to determine number of fertile and infertile eggs. After candling the fertile eggs of each replicated tray were transferred to separate hatching trays and placed in a hatcher. The hatcher was operated at a temperature 98°F and a relative humidity of 65-70%. After 21 days of incubation all hatched chicks were taken out of hatcher and shifted to chick room of hatchery. The hatchability parameters were studied during and after incubation period like fertility, hatchability, dead germ, dead in shell, moisture loss, shell weight and body weight of day-old chicks were recorded. Ten eggs from each group were selected randomly and subjected to study

the external and internal egg quality traits. Statistical analysis was performed using one-way analysis of variance (ANOVA) to compare group means, and the significance of differences among treatments was tested at the 5% probability level using IBM SPSS Statistics (Version 30.0; IBM Corp., Armonk, NY, USA, 2023).

RESULTS AND DISCUSSION

External egg quality traits

The average egg weight (g) was 42.16±0.34, 48.09±0.55, 52.65±0.47, 57.98±0.41 and 63.34±0.68 respectively for T₁, T₂, T₃, T₄ and T₅ groups which differed significantly (P<0.05). The egg shape index (%) ranged from 74.54±1.74 to 76.69±1.65 which are within the normal range of optimum shape of egg. Similar shape index was also reported by Sheikh *et al.* (2018) in indigenous chicken eggs, Jaishankar *et al.* (2020) in Kadaknath eggs (73.64%), Kumar *et al.* (2022) in Aseel (74.75±0.47%) and Kadaknath eggs (74.02±0.46%), and Suresh *et al.* (2025) in Vanaraja chicken eggs. The lower values of shape index in Vanaraja eggs were reported by Sheikh *et al.* (2018). The egg specific gravity (mg/dL) ranged from 1.061±0.06 in T₂ to 1.092±0.03 in T₅ group. The egg shell colour score was same for all the groups (2.00).

Internal egg quality traits

Shell thickness (mm) ranged from 0.281±0.02 in T₁ to 0.308±0.01 in T₅ group. Slightly higher shell thickness was reported by Suresh *et al.* (2025). Shell weight (g) ranged from 4.53±0.31 in T₁ to 6.81±0.51 in T₅ group with significant (P≤0.05) differences among the groups. Similar result regarding shell weight was also reported by John-Jaja *et al.* (2016). Albumen diameter (mm) ranged from 85.56±3.75 in T₁ to 93.38±5.02 in T₅ group. Significantly (P≤0.05) highest albumen weight (g) was recorded in T₅ (38.59±1.12g) and lowest in T₁ (24.31±1.87g) group. Yolk diameter (mm) and yolk height (mm) ranged from 42.24±1.66 in T₁ to 46.32±2.03 in T₃ and 15.91±2.45 in T₂ to 18.08±1.71 in T₅ group, respectively. Yolk weight (g) ranged from 13.30±0.51g in T₁ to 17.91±0.65g in T₅ group. Yolk diameter, yolk height and yolk weight differed

significantly ($P \leq 0.05$) among the groups. Yolk index ranged from 0.353 ± 0.05 to 0.404 ± 0.07 which is lower than reported by Sheikh *et al.* (2018) in Vanaraja eggs. The Haugh unit (HU) score is the measure of albumen quality which determines the quality of the egg, and it ranged from 69.87 ± 4.03 in T_1 to 73.09 ± 3.21 in T_3 group with significant ($P \leq 0.05$) differences among the groups. Similar results were also reported by Sheikh *et al.* (2018) in crossbred chicken eggs. However, higher values of HU score was reported by Jaishankar *et al.* (2020) in Kadaknath eggs (87.28 ± 1.85), Supasini *et al.* (2024) in indigenous chicken eggs in Srilanka, Suresh *et al.* (2025) in Vanaraja eggs (84.88 ± 0.49) and lower HU score was reported by Gautam *et al.* (2021) in

large sized Vanaraja eggs (58.91 ± 3.63). Shell thickness, albumen height (mm), albumen index and yolk index did not differ significantly ($P > 0.05$) among the groups. Ukwu *et al.* (2017) also found that yolk index did not differ significantly ($P > 0.05$), with different egg weights.

Moisture loss, fertility and hatchability

During the incubation process, moisture loss creates an air cell, essential for the growing embryo to breathe. The moisture loss (%) was recorded to be ranged between 10.08 ± 0.02 in T_1 and 11.46 ± 0.15 in T_5 groups, as size of egg increased the moisture loss also increased without any significant ($P > 0.05$) difference which might be due to diffusion of water

Table 1: External egg quality traits of Vanaraja chicken eggs of different sizes

Parameters	T_1 (40-45g)	T_2 (45-50g)	T_3 (50-55g)	T_4 (55-60g)	T_5 (60-65 g)
Egg weight (g)	$42.16^a \pm 0.34$	$48.09^b \pm 0.55$	$52.65^c \pm 0.47$	$57.98^d \pm 0.41$	$63.34^e \pm 0.68$
Egg shape index (%)	76.69 ± 1.65	75.33 ± 2.72	75.06 ± 1.97	74.54 ± 1.74	74.81 ± 1.83
Egg specific gravity (mg/dL)	1.073 ± 0.02	1.061 ± 0.06	1.084 ± 0.01	1.076 ± 0.02	1.092 ± 0.03
Egg shell colour score	2.00	2.00	2.00	2.00	2.00

Means across columns in a same row bearing different superscript differ significantly ($P \leq 0.05$)

Table 2: Internal egg quality traits of Vanaraja chicken eggs of different sizes

Parameters	T_1 (40-45g)	T_2 (45-50g)	T_3 (50-55g)	T_4 (55-60g)	T_5 (60-65 g)
Shell thickness (mm)	0.281 ± 0.02	0.292 ± 0.01	0.302 ± 0.02	0.295 ± 0.01	0.308 ± 0.01
Shell weight (g)	$4.53^a \pm 0.31$	$4.94^a \pm 0.29$	$5.39^a \pm 0.40$	$5.68^{ab} \pm 0.38$	$6.81^b \pm 0.51$
Albumen diameter (mm)	$85.56^a \pm 3.75$	$87.34^{ab} \pm 2.96$	$90.74^b \pm 5.61$	$89.90^b \pm 4.08$	$93.38^c \pm 5.02$
Albumen height (mm)	2.82 ± 0.21	2.75 ± 0.19	3.15 ± 0.27	2.99 ± 0.16	3.24 ± 0.25
Albumen index	0.033 ± 0.004	0.032 ± 0.003	0.034 ± 0.003	0.033 ± 0.002	0.034 ± 0.003
Albumen weight (g)	$24.31^a \pm 0.87$	$28.12^{ab} \pm 0.58$	$31.57^{bc} \pm 0.94$	$35.0^c \pm 1.08$	$38.59^c \pm 1.12$
Yolk diameter (mm)	$42.24^a \pm 1.66$	$43.97^{ab} \pm 1.89$	$46.32^b \pm 2.03$	$45.68^b \pm 1.80$	$44.73^b \pm 2.16$
Yolk height (mm)	$16.45^a \pm 2.52$	$15.91^a \pm 2.45$	$17.20^a \pm 2.77$	$16.12^a \pm 1.60$	$18.08^b \pm 1.71$
Yolk index	0.389 ± 0.06	0.361 ± 0.05	0.371 ± 0.04	0.353 ± 0.05	0.404 ± 0.07
Yolk Weight (g)	$13.30^a \pm 0.54$	$15.00^a \pm 0.66$	$15.68^{ab} \pm 0.42$	$17.26^b \pm 0.81$	$17.91^b \pm 0.75$
Haugh unit scores	$69.87^a \pm 4.03$	$70.62^a \pm 3.65$	$73.09^b \pm 3.21$	$72.51^b \pm 3.01$	$70.11^a \pm 2.76$

Means across columns in a same row bearing different small case superscript differ significantly ($P \leq 0.05$)

Table 3: Effect of Egg size on hatching performance and Day-Old Chick Weight in Vanaraja chicken

Parameters	T_1 (40-45g)	T_2 (45-50g)	T_3 (50-55g)	T_4 (55-60g)	T_5 (60-65 g)
Average Egg weight (g)	$43.51^a \pm 1.27$	$47.62^b \pm 1.37$	$52.53^c \pm 2.03$	$57.62^d \pm 1.58$	$62.53^e \pm 1.86$
Moisture loss (%) during incubation	10.08 ± 0.02	10.56 ± 0.29	10.95 ± 0.25	11.02 ± 0.10	11.46 ± 0.15
Fertility (%)	$71.30^a \pm 4.67$	$78.00^b \pm 8.00$	$83.50^c \pm 8.50$	$84.00^c \pm 7.02$	$81.00^c \pm 3.10$
Hatchability TES (%)	$49.00^a \pm 4.10$	$58.00^b \pm 6.31$	$65.50^c \pm 4.85$	$64.50^c \pm 5.50$	$54.70^b \pm 2.75$
Hatchability FES (%)	$68.00^a \pm 8.00$	$73.90^b \pm 4.76$	$76.70^b \pm 3.68$	$76.40^b \pm 3.74$	$66.50^a \pm 3.15$
Dead germ (%)	$7.33^c \pm 0.67$	3.50 ± 0.49^a	3.57 ± 0.53^a	4.50 ± 0.58^{ab}	5.75 ± 1.05^b
Dead in Shell (%)	$19.25^b \pm 1.15$	$18.50^b \pm 1.58$	$14.50^a \pm 1.50$	$15.00^a \pm 1.23$	$18.70^b \pm 4.25$
DOC weight (g)	$27.67^a \pm 1.49$	$30.96^{ab} \pm 1.20$	$34.65^b \pm 0.77$	$38.00^c \pm 1.40$	$42.34^d \pm 0.85$

Means across columns in a same row bearing different small case superscript differ significantly ($P \leq 0.05$)

through the shell (Tona *et al.*, 2001). The egg white contains proportionately almost double the amount of water compared to yolk and large eggs have a greater proportion of albumen thus greater moisture content than small eggs (Abanikannda *et al.*, 2011). Yousaf *et al.* (2024) reported that total egg water loss of ~12% of initial fresh mass during incubation results in highest hatching success. The fertility (%) ranged from 71.30 ± 4.67 in T_1 to 84.00 ± 7.02 in T_4 group. Hatchability on TES (%) ranged from 49.00 ± 9.00 in T_1 to 65.50 ± 8.50 in T_3 group and on FES (%) was from 66.50 ± 3.15 in T_5 to 76.70 ± 3.68 in T_3 group. The fertility and hatchability were better in T_3 and T_4 groups with significant ($P \leq 0.05$) differences and this result is in agreement with the findings of Nowaczewski *et al.* (2022) who also observed that the eggs of intermediate size range would hatch better. However, Patel *et al.* (2018) reported higher fertility ($86.31 \pm 0.23\%$) and higher hatchability (%) on TES (64.18 ± 0.71) and FES (74.36 ± 0.63) in Vanaraja eggs and Jaishankar *et al.* (2020) reported higher fertility ($91.11 \pm 0.77\%$) and higher hatchability on TES ($73.89 \pm 1.43\%$) and FES ($80.95 \pm 1.19\%$) in Kadaknath chicken eggs. These differences might be due to different environmental and managerial conditions. The dead-germ (%) was significantly (≤ 0.05) highest in T_1 (7.33 ± 0.67) and lowest in T_2 and T_3 (3.50 ± 0.49). Nowaczewski *et al.* (2022) and Kumar *et al.* (2024) also reported that the embryonic mortality was highest in large-sized and very small sized eggs. Dead-in-shell (%) ranged from 14.50 ± 0.50 in T_3 to 21.70 ± 4.25 in T_5 group with significant ($P \leq 0.05$) differences among the groups. Mabothea *et al.* (2025) also observed similar dead-in-shell rates in broiler breeders. The day-old chick weight (g) was significantly ($P \leq 0.05$) highest in T_5 (42.34 ± 0.85) and lowest in T_1 (27.67 ± 1.49) group. A significant ($P \leq 0.05$) linear increase in day old chick body weight was observed with increasing egg weight which might be due to larger eggs contained more nutrients than small eggs so that developing embryos from heavier eggs tended to have more nutrients for their growth requirements during incubation, hence heavy chicks hatched from large eggs (Williams, 1994). Chicks hatched from smaller eggs exhibited lower body weight compared to those originating from larger eggs; similar finding

indicating a strong relationship between egg weight and chick hatch weight were reported by Ewonetu and Kasaye (2018).

CONCLUSION

The present study revealed that egg size is a significant factor in determining egg quality traits, hatchability performance, embryonic mortality, and day-old chick weight in Vanaraja chicken. Medium sized eggs (50-60g) had better external and internal quality attributes, along with higher fertility and hatchability percentages and lower dead-in-shell rates compared to smaller and larger eggs. Day-old chick weight had a positive linear relationship with egg weight. The chicks with higher body weight at hatch grows faster than their counterpart of lower body weight which is an important factor in poultry farming. Based on the results, it can be concluded that the optimal weight range of 50-60g is the best choice for better hatchery performance in Vanaraja chicken.

REFERENCES

- Abanikannda, O.T.F., Leigh, A.O. and Giwa, A.O. 2011. Influence of egg weight, breed, and age of hens on weight loss of hatching broiler eggs. *Archiva Zootechnica*, 14: 343-356.
- Agbehadzi, R. K., Meteyake, H. T., Adjei-Mensah, B., Sasu, P., Koranteng, A. A., Dassidi, N., Hamidu, J. A. and Tona, K. (2024). Effects of layer breeder age and reduced incubator oxygen concentrations on embryo development, hatching events, chick quality, embryonic mortality and hatchability of fertile eggs. *Heliyon*, 10(22): e40468.
- Ewonetu, K. S. and Kasaye, A. (2018). Effect of egg weight on post-hatch performance of White Leghorn chicken breed from day-old to laying age. *Journal of Poultry Research*, 15(2): 16-22.
- Fathi, M., Abou-Emera, O., Al-Homidan, I., Galal, A. and Rayan, G. (2022). Effect of genotype and egg weight on hatchability properties and embryonic mortality pattern of native

- chicken populations. *Poultry Science*, 101(11): 102129.
- Gandhimathi, D., Murugan, M. and Edwin, S. C. (2020). Effect of genetic makeup and season on fertility and hatchability of chicken varieties of rural farming. *Indian Journal of Veterinary and Animal Sciences Research*, 49(6): 55–59.
- Gautam, A. Khan, A. A., Hamadani, H., Haq, Z. and Sofi, A. H. (2021). Egg quality traits of different sized hatching eggs from Vanaraja parent stock reared in Kashmir. *SKUAST Journal of Research*, 23(2): 212-216.
- Jaishankar, S., R. Jyothi Priya, A. Sheeba and Ilavarasan, S. (2020). Productive and Reproductive Performance of Kadaknath Chicken under Semi-intensive System. *International Journal of Current Microbiology and Applied Science*, 9(4): 513-517.
- John-Jaja, S. A., Udoh, U. H. and Nwokolo, S. C. (2016). Repeatability estimates of egg weight and egg-shell weight under various production periods for Bovan Nera Black laying chickens. *Beni-Suef University Journal of Basic and Applied Sciences*, 5(4): 389-394.
- Kuda, N. T. (2023). Factors influencing fertility and hatchability of poultry. *Journal of Biology, Agriculture and Healthcare*, 13(8): 1–7.
- Kumar, M., Dahiya, S.P., Ratwan, P., Sheoran, N., Kumar, S. and Kumar, N. (2022). Assessment of egg quality and biochemical parameters of Aseel and Kadaknath indigenous chicken breeds of India under backyard poultry farming. *Poultry Science*, 101(2): 101589
- Kumar, R., Jamwal, K., Bhardwaj, N., Sankhyan, V., Dinesh, K. and Katich's (2025). Effects of egg size on hatchability parameters and chick quality in native chicken: Egg size on hatchability. *Indian Journal of Poultry Science*, 59(2):193-198.
- Mabotha, T. R., Nephawe, K. A. and Mpofo, T. J. (2025). Effect of flock age, egg size, and incubator type on embryo development and hatchability in the Indian River Broiler breeders in South Africa: Embryo development in Indian River Broiler breeder. *Letters In Animal Biology*, 5(2): 136–145.
- Mbajjorgu C.A. and Ramaphala N.O. (2026). Insight into egg weight and its impact on chick hatch-weight, hatchability and subsequent growth indices in chickens- a review. *Indian Journal of Animal Research*, 48(3): 209-213.
- Nowaczewski, S., Babuszkiewicz, M., Szablewski, T., Stuper-Szablewska, K., Cegielska-Radziejewska, R., Tomczyk, L., Kaczmarek, S., Sechman, A., Lis, M. W., Kwa[niewska, M., Racewicz, P., Jarosz, A., Ciszewski, A., Nowak, T. and Hejdysz, M. (2022). Effect of weight and storage time of broiler breeders' eggs on morphology and biochemical features of eggs, embryogenesis, hatchability, and chick quality. *Animal: An International Journal of Animal Bioscience*, 16(7): 100564.
- Patel, N., Sundi, B., Prasad, S., Kumar, R. and Man, B. (2018). Reproductive and egg quality traits of Vanaraja birds under different system of management. *International Journal of Current Microbiology and Applied Sciences, Special Issue-7*: 696–699.
- Peric L., Mitrakovic, M., Tomic, B., Orehovacki, I. and Meijerhof, R. (2022). Effects of flock age, place of oviposition and cleaning treatments of hatching eggs on hatchability in broiler breeders. *Journal of Applied Poultry Research*, 31: 100279.
- Rathnayake, C., Herath, S., Pathirana, S., Lakmini, G. and Thilakarathne, S. (2024). Development of a cost-effective and high-efficiency incubator for small-scale poultry farmers to maximize hatchability and brood quality. *Indian Journal of Animal Production and Management*, 40(4): 243–250.
- Sheikh, I. U., Kalita, N., Mahanta, J. D. and Hussain, J. (2018). Studies on certain egg qualities of indigenous, Vanaraja, and crossbred (PB2 × indigenous) chickens under intensive and

- backyard systems of rearing. *International Journal of Current Microbiology and Applied Sciences*, 7(4): 1690–1695.
- Supasini, K., Akram, A.T.A., Ahamed, A.S., Musthafa, M.M. and Marikar, F.M.M.T. (2024). Evaluating the relationship between egg weight and village chicken egg quality in Sri Lanka. *Journal of Natural and Applied Sciences Pakistan*, 6(1):1663-1673.
- Suresh, K., Sreenivas, D., Jaya Laxmi, P., Raj Kumar, U. and Pragnya, Ch. (2025). Study of egg quality traits in Vanaraja male line (PD-1). *International Journal of Advanced Biochemistry Research*, SP-9(7): 764-766.
- Tona, K., Bamelis, F., Couke, W., Bruggeman, V. and Decuyper, E. (2001). Relationship between broiler breeders age and egg weight loss and embryonic mortality during incubation in large scale conditions. *Journal of Applied Poultry Research*, 10: 221-227.
- Ukwu, H.O., Ezihe, C.O., Asaa, S.K. and Anyogo, M.E. (2017). Effect of egg weight on external and internal egg quality traits of Isa Brown egg layer chickens in Nigeria. *Journal of Animal Science and Veterinary Medicine*, 2(4): 126-32.
- Williams, T.D. (1994). Intraspecific variation in egg size and egg composition in birds: effects on offspring fitness. *Biological Reviews*, 69: 35-59.
- Yousaf, A., Yasoob, T.B., Bilal, M., Anwar, D.M., Ghaffar, A., Lodhi, S.M.K. and Malik, M. (2024). Effect of Egg Weight on Water Loss and Chicks Yield of Broiler Breeders. *Biomedical Journal of Science and Technology Research*, 56(2). BJSTR. MS.ID.008822.

Received: February 23, 2026

Accepted: March 25, 2026